

Nonoperative Management of Cervical Radiculopathy

MARC A. CHILDRESS, MD, *Fairfax Family Practice Primary Care Sports Medicine Fellowship, Fairfax, Virginia*

BLAIR A. BECKER, MD, *Group Health Family Medicine Residency, Seattle, Washington*

Cervical radiculopathy describes pain in one or both of the upper extremities, often in the setting of neck pain, secondary to compression or irritation of nerve roots in the cervical spine. It can be accompanied by motor, sensory, or reflex deficits and is most prevalent in persons 50 to 54 years of age. Cervical radiculopathy most often stems from degenerative disease in the cervical spine. The most common examination findings are painful neck movements and muscle spasm. Diminished deep tendon reflexes, particularly of the triceps, are the most common neurologic finding. The Spurling test, shoulder abduction test, and upper limb tension test can be used to confirm the diagnosis. Imaging is not required unless there is a history of trauma, persistent symptoms, or red flags for malignancy, myelopathy, or abscess. Electrodiagnostic testing is not needed if the diagnosis is clear, but has clinical utility when peripheral neuropathy of the upper extremity is a likely alternate diagnosis. Patients should be reassured that most cases will resolve regardless of the type of treatment. Nonoperative treatment includes physical therapy involving strengthening, stretching, and potentially traction, as well as nonsteroidal anti-inflammatory drugs, muscle relaxants, and massage. Epidural steroid injections may be helpful but have higher risks of serious complications. In patients with red flag symptoms or persistent symptoms after four to six weeks of treatment, magnetic resonance imaging can identify pathology amenable to epidural steroid injections or surgery. (*Am Fam Physician*. 2016;93(9):746-754. Copyright © 2016 American Academy of Family Physicians.)

CME This clinical content conforms to AAFP criteria for continuing medical education (CME). See CME Quiz Questions on page 736.

Author disclosure: No relevant financial affiliations.

► **Patient information:** A handout on this topic, written by the authors of this article, is available at <http://www.aafp.org/aafp/2016/0501/p746-s1.html>.

Scan the QR code below with your mobile device for easy access to the patient information handout on the AAFP mobile site.



Cervical radiculopathy describes pain in one or both of the upper extremities, often in the setting of neck pain, secondary to compression or irritation of nerve roots in the cervical spine. It can be accompanied by motor, sensory, or reflex deficits.¹ The annual incidence is 107 per 100,000 men, and 64 per 100,000 women. It is most prevalent in persons 50 to 54 years of age.²

The underlying pathology is typically degenerative, including spondylosis and disk herniation.² Osteoarthritis in these conditions causes bony hypertrophy, most commonly at the facet joints or uncovertebral joints.³ In 22% of cases, nerve compression occurs as a result of disk herniation.² Nerve impingement occurs more rarely with trauma.

Anatomy

The cervical nerve roots exit the spine laterally through neural foramina. Each foramen is bordered by joints that are prone to degenerative arthritis, or spondylosis (*Figure 1*). Anterior to the foramen is the uncovertebral

joint, and posterior to the foramen is the facet joint.

The intervertebral disk lies anterior and medial to the neural foramen, and in the setting of herniation, can protrude onto the exiting nerve root. Trauma can lead to instability that injures the nerve root proximally as well.³

In a feature unique to the cervical spine, each nerve root from C1 to C7 exits above its corresponding vertebral level. The exception is the C8 nerve root, which exits below the seventh vertebra. The motor portion of the cervical nerves (ventral nerve root) exits the spinal cord anteriorly, whereas the sensory portion (dorsal nerve root) exits posteriorly before forming the dorsal root ganglion and joining the ventral nerve root to form a mixed spinal nerve.

SPONDYLOSIS TYPES

Spondylosis leading to radiculopathy can occur at the uncovertebral and facet joints. The uncovertebral joint's location anterior to the nerve root means that bony hypertrophy here tends to affect the anterior aspect of the nerve root. Conversely, facet joint arthritis




Figure 1. Cervical spine anatomy. Cervical nerve root impingement commonly results from bony hypertrophy at the uncovertebral joint or the facet joint, or from disk herniation.

affects the posterior aspect of the nerve root. Degenerative loss of disk height and resulting arthritic hypertrophy can also decrease the diameter of the neural foramen and can contribute to nerve root impingement.³

DISK HERNIATION TYPES

There are three main types of disk herniation that lead to nerve root impingement (*Figure 2*). The most common is intraforaminal, which results in predominantly sensory radicular symptoms. The next most common is posterolateral, which results in weakness and potentially muscle atrophy. Rarer midline herniations directly compress the spinal cord and result in symptoms of myelopathy, such as upper extremity numbness, weakness, gait disturbance, ataxia, and urinary incontinence.⁴

History

Radicular pain is the most common symptom in cervical radiculopathy, followed by paresthesia. Weakness is reported by about 15% of patients.² The distribution of numbness should be noted. In most cases, the affected nerve root can be identified by the history and physical examination alone⁵ (*Figure 3*). Some patients also report pain that radiates to the posterior shoulder and periscapular region, which is relatively nonspecific.³ Patients may report worsening of symptoms with neck extension or lateral flexion to the affected side.

The patient should be asked about red flag signs and symptoms of myelopathy, malignancy, and spinal abscess that require immediate workup (*Table 1*).

Physical Examination

The most common physical examination findings in persons with cervical radiculopathy are painful neck movements and muscle spasm. Diminished deep tendon reflex is the most common objective neurologic finding, with triceps involvement being the most prevalent.




Figure 2. Disk herniation types. (A) Intraforaminal. (B) Posterolateral. (C) Midline.

BEST PRACTICES IN NEUROLOGY: RECOMMENDATIONS FROM THE CHOOSING WISELY CAMPAIGN

Recommendation	Sponsoring organization
Do not do nerve conduction studies without also doing needle electromyography to test for radiculopathy, a pinched nerve in the neck or back.	American Association of Neuromuscular & Electrodiagnostic Medicine

Source: For more information on the Choosing Wisely Campaign, see <http://www.choosingwisely.org>. For supporting citations and to search Choosing Wisely recommendations relevant to primary care, see <http://www.aafp.org/afp/recommendations/search.htm>.




Figure 3. Relevant cervical dermatomes. Sensory symptoms often correlate with specific dermatomes.

Table 1. Red Flags in the Evaluation of Radicular Neck Pain

Condition	Historic findings	Examination findings
Malignancy	Fever, history of cancer, night pain, weight loss	Variable, gait abnormality; unilateral neurologic findings more common than bilateral
Myelopathy	Decreased dexterity, urinary urgency	Ataxia, clonus, hyperreflexia
Spinal abscess	Fever, history of intravenous drug use, immunocompromise	Neurologic deficit

Table 2. Neurologic Patterns of Cervical Nerve Roots

Nerve root	Interspace	Pain distribution	Abnormalities		
			Motor	Sensory	Reflex
C4	C3 to C4	Lower neck, trapezius	—	Cape distribution (i.e., lower neck and upper shoulder girdle)	—
C5	C4 to C5	Lateral arm, neck, shoulder	Deltoid, elbow flexion	Lateral arm	Biceps
C6	C5 to C6	Dorsolateral arm, neck, thumb	Biceps, wrist extension	Lateral forearm, thumb	Brachioradialis
C7	C6 to C7	Dorsolateral forearm, middle finger, neck	Triceps, wrist flexion	Dorsal forearm, long finger	Triceps
C8	C7 to T1	Medial forearm, neck, ulnar digits	Finger flexors	Medial forearm, ulnar digits	—
T1	T1-T2	Ulnar forearm	Finger intrinsics	Ulnar forearm	—

Adapted with permission from Eubanks JD. Cervical radiculopathy: nonoperative management of neck pain and radicular symptoms. *Am Fam Physician.* 2010;81(1):35.

Weakness is the next most common finding.² The most common nerve root affected is C7, followed by C6.^{2,6,7}

Knowledge of cervical myotomes and dermatomes is helpful (Table 2⁸), but radicular pain may manifest outside classic dermatomal borders.⁴

PROVOCATIVE MANEUVERS

The Spurling test is a highly specific^{7,9-11} and sensitive^{11,12} maneuver validated by operative, magnetic resonance imaging (MRI), and electrodiagnostic findings. It involves passively moving the patient's neck into lateral flexion and extension, then applying gentle downward axial compression (Figure 4). The purpose of this maneuver is to constrict the neural foramen; a positive result is the reproduction of radicular symptoms.

The shoulder abduction test is similar in specificity to the Spurling test, based on electrodiagnostic correlation.⁷ The test involves placing the palm of the affected limb on top of the patient's head. A positive result is the relief of radicular symptoms.

The more complicated upper limb tension test is less commonly used, but is the most sensitive test for ruling out cervical radiculopathy^{7,10,12} (Figure 5). The maneuver can be thought of as a cervical nerve tension analog to the straight leg raise test for the lumbar spine. It consists of initially placing the patient in a supine position with the shoulder in a neutral position and the elbow and wrist flexed. The examiner then abducts the shoulder to



ILLUSTRATION BY MARCIA HARTSOCK

Figure 4. Spurling maneuver. Lateral flexion and extension of the neck with axial compression.



Figure 5. Upper limb tension test. (A) Scapular depression with shoulder abduction. (B) Contralateral flexion of the neck with extension of the elbow, wrist, and fingers, and supination of the wrist.

90 degrees, extends the elbow and fingers, and extends and supinates the wrist while the patient deviates the neck to the contralateral and then ipsilateral sides. A positive result is reproduction of pain at any step of the maneuver.

There have been reports that the Valsalva maneuver may provoke radicular symptoms, but the sensitivity and specificity of this test are not known.³

Other Causes of Cervical Nerve Root Impingement

Other causes of nerve root impingement often present with a unique constellation of symptoms (*Table 3*). Spinal tumors most often cause myelopathy, although osteochondromas and schwannomas can cause radiculopathy. Because these lesions tend to occur within the dura mater, the Valsalva maneuver can exacerbate symptoms. Tumors stemming from thyroid, esophageal, pharyngeal, and lung tissue have been reported to compress individual cervical nerves distal to the neural foramen, as have sarcoidosis and arteriovenous malformations.³

**Imaging
RADIOGRAPHY**

Although plain radiographs of the cervical spine are useful for ruling out instability (*Figure 6*), they are relatively nonspecific for diagnosing cervical radiculopathy. About 65% of asymptomatic patients 50 to 59 years of age will have radiographic evidence of significant cervical spine degeneration, regardless of radiculopathy symptoms.¹³ C5 to C6 is the most common level affected, followed closely by C6 to C7. Plain radiography should be ordered if there is a history of trauma, red flags for malignancy, or failure to improve at four to six weeks. The series should include an oblique view, which can sometimes reveal bony hypertrophy at the neural foramina.

MRI

MRI is indicated in patients with complex cervical radiculopathy (*Figure 7*), which is defined by a high suspicion for myelopathy or abscess, persistent or progressive objective neurologic findings, or failure to improve after four to six weeks of conservative treatment.^{1,14-18} MRI is not helpful in most cases of cervical radiculopathy because of its significant rates of false-negative and false-positive findings. About 57% of patients older than 64 years who do not have symptoms of cervical radiculopathy have evidence of disk herniation, and 26% have spinal cord impingement.¹⁹

COMPUTED TOMOGRAPHIC MYELOGRAPHY

MRI has mostly supplanted computed tomographic myelography as the first-line imaging modality for

Table 3. Differential Diagnosis for Cervical Radiculopathy

Condition	Key clinical feature	Diagnostic evaluation
Abscess	Fever, neurologic deficit, pain	Blood cultures, complete blood count, erythrocyte sedimentation rate, MRI
Anterior interosseus nerve entrapment	Grip and pinch weakness, no pain	EMG
Arteriovenous malformation	Numbness, paresthasias, variable pain, weakness	MRI, ultrasonography
Carpal tunnel syndrome	Numbness in radial 3½ fingers, paresthasias, thenar weakness	Hand symptom diagram, Tinel and Phalen tests with or without EMG
Cervical myelopathy	Ataxia, decreased dexterity, urinary urgency	MRI, tests for clonus and hyperreflexia
Cubital tunnel syndrome	Flexor digitorum profundus weakness, numbness in ulnar half of ring and little fingers, paresthasias	Tinel test with or without EMG
Extraspinal malignancy (e.g., thyroid, esophageal, pharyngeal, or Pancoast tumor)	Variable	Variable
Herpes zoster	Vesicular rash	Viral culture
Parsonage-Turner syndrome (brachial plexopathy)	Acute onset of pain, then numbness and weakness	EMG with or without MRI
Posterior interosseus nerve entrapment	Finger and wrist weakness, pain	EMG with or without MRI
Radial tunnel syndrome	Pain only at the radial forearm	Diagnostic injection
Reflex sympathetic dystrophy	Edema, pain, skin discoloration	Bone scintigraphy
Rotator cuff tendinosis	Shoulder pain with potential radiation to arm	Ultrasonography
Thoracic outlet syndrome	Pain, swelling, vascular insufficiency	Adson, Roos, and Wright tests with or without angiography
Tumor	Fever, gait abnormality, pain, paresthasias, weakness, weight loss	MRI

EMG = electromyography; MRI = magnetic resonance imaging.

complex cervical radiculopathy.¹⁴ However, computed tomographic myelography remains the test of choice to clarify clinically apparent nerve root impingement in patients with equivocal MRI findings.^{14,20}

Electrodiagnostic Testing

There is insufficient evidence to support the routine use of electrodiagnostic testing in the workup of cervical radiculopathy.¹ Electromyography does, however, have clinical utility when peripheral neuropathy of the upper extremity is a likely alternate diagnosis. It can be challenging to differentiate proximal from distal nerve root impingement, but a working knowledge of common peripheral neuropathies is useful (*Table 4*).

Nonoperative Treatment

Most patients with cervical radiculopathy will improve with nonoperative care^{2,15} (*Figure 8*). However, no treatment modalities are supported by evidence from high-quality studies.



Figure 6. Cervical radiography, oblique view. Bony intervertebral foraminal stenosis at the left C3-C4 level caused by uncovertebral osteophyte and facet joint hypertrophy.




Figure 7. T2-weighted cervical magnetic resonance imaging. (A) Sagittal view of a large paracentral/foraminal disk protrusion at the C4-C5 level indenting the anterior thecal sac and flattening the anterior surface of the cord. (B) Axial view of the same lesion. The disk protrusion caused severe left intervertebral foraminal stenosis and compressed the left C5 nerve within the foramen.

Table 4. Differentiating Common Peripheral and Cervical Nerve Root Compressions

Condition	Affected nerve	Signs and symptoms	Cervical nerve root mimicker	Distinguishing feature
Carpal tunnel syndrome	Median	Paresthesias, thenar weakness, numbness in radial 3½ fingers	C6, C7	No triceps or wrist extensor weakness in carpal tunnel syndrome
Cubital tunnel syndrome	Ulnar	Paresthesias, grasp weakness, numbness in ulnar half of ring and little fingers	C8, T1	No thumb interphalangeal flexion weakness in cubital tunnel syndrome
Posterior interosseus nerve entrapment	Peripheral interosseus nerve branch of the radial nerve	Pain, wrist and finger extensor weakness	C7	No triceps or wrist flexor weakness in posterior interosseus nerve entrapment

PHYSICAL THERAPY

A Cochrane review found low-quality evidence to support cervical, shoulder, scapulothoracic, and upper arm strengthening and stretching in the acute phase for treatment of radicular pain.^{16,17} A randomized controlled trial (RCT) showed significant improvement in patients receiving twice-weekly supervised physical therapy and home physical therapy compared with control patients in the first six weeks of cervical radiculopathy.¹⁸ Several cohort studies and randomized trials drew similar conclusions.^{21,22}

TRACTION

Many physical therapy regimens for cervical radiculopathy include mechanical traction. Based on low-quality evidence, several systematic reviews and RCTs have concluded that traction is no better than placebo.²³⁻²⁸ A more recent RCT found that 10 sessions of supervised

mechanical traction over four weeks, in addition to physical therapy, were superior to physical therapy alone at six and 12 months.²⁹ There was also significant improvement at 12 months in a group treated with four weeks of physical therapy and home over-the-door traction. Two other studies reached similar conclusions.^{30,31}

SOFT COLLAR

Although one study found superior gains in function with a semi-hard collar compared with physical therapy and placebo, a systematic review and several RCTs found little improvement in pain with the use of soft and semi-hard collars for cervical radiculopathy.^{24,27,31}

MASSAGE

A Cochrane review of 15 trials that included studies on neck pain with and without radiculopathy found low-quality evidence to support the use of classical, modified

Management of Cervical Radiculopathy




Figure 8. Algorithm for the management of cervical radiculopathy. (MRI = magnetic resonance imaging.)

strain/counterstrain, and traditional Chinese massages for the improvement of pain and function compared with placebo or education alone.³² Further studies are needed to establish the benefit of these treatments.

ORAL MEDICATIONS

Nonsteroidal anti-inflammatory drugs and muscle relaxants are often prescribed for treatment of acute cervical radiculopathy. A Cochrane review concluded that there is limited evidence to support the use of these agents in

the setting of acute neck pain, but did not specifically address radiculopathy.³³ A small RCT found that a short course of oral corticosteroids reduced radiculopathy-related pain in the short term.³⁴

STERIOD INJECTION

Steroid injection can be considered for patients whose symptoms persist after four to six weeks of conservative management. A 2007 Cochrane review found low-quality evidence to support the use of epidural

SORT: KEY RECOMMENDATIONS FOR PRACTICE

Clinical recommendation	Evidence rating	References
Cervical radiculopathy can be diagnosed with the patient history and physical examination alone.	C	1, 5, 6
Diagnostic sensitivity and specificity is improved with the use of the Spurling and upper limb tension tests.	C	7, 10-12
Magnetic resonance imaging is the preferred imaging modality in patients who have no improvement after four to six weeks of nonoperative treatment or progression of objective neurologic deficit.	C	1, 14-18
Physical therapy with strengthening and stretching can be beneficial in acute cervical radiculopathy.	A	16, 22
Most patients with cervical radiculopathy will improve regardless of nonoperative treatment modality.	C	1, 2, 21
Surgical referral or epidural steroid injections should be considered if there is no improvement after four to eight weeks of nonoperative treatment.	C	1, 38, 42

A = consistent, good-quality patient-oriented evidence; B = inconsistent or limited-quality patient-oriented evidence; C = consensus, disease-oriented evidence, usual practice, expert opinion, or case series. For information about the SORT evidence rating system, go to <http://www.aafp.org/afpsort>.

steroid injections in patients with chronic cervical radiculopathy.³⁵ A more recent systematic review found good-quality evidence to support steroid injections for cervical radiculopathy caused by disk herniation, but only fair evidence for radiculopathy caused by spondylosis.³⁶ An RCT showed significant benefit in pain and function with epidural steroid injections for at least one year in patients who had not improved with physical therapy and nonsteroidal anti-inflammatory drugs.³⁷ Several cohort studies showed significant improvements in recalcitrant radicular pain with epidural steroid injections, in most cases for at least one year.^{9,35,38-40} Patients should be counseled about the risk of potential complications, including dural puncture, meningitis, epidural abscess, and nerve root injury.

Prognosis

Most patients with cervical radiculopathy will improve regardless of treatment modality.² In fact, roughly 88% will improve within four weeks of nonoperative management.¹⁵ In a retrospective case series, 80% of those with objective weakness or reflex deficit improved within three weeks of conservative management.⁴¹ Repeat examination at follow-up is crucial. Progression of an objective neurologic finding at any point may signify advancing nerve root compression and should trigger an MRI and referral to a spinal surgeon.

The optimal timing for referral in cases of recalcitrant but nonprogressive radiculopathy is not clear. There is evidence to support referral within four to eight weeks.^{15,42} Radiography and MRI can be considered if there is no improvement at four to six weeks. If imaging reveals evidence of nerve root impingement that correlates with physical examination findings, referral to a spinal surgeon is recommended.

Data Sources: The terms cervical radiculopathy and non operative and cervical radiculopathy were searched through PubMed, Cochrane, Essential Evidence Plus, and the Agency for Healthcare Research and Quality National Guideline Clearinghouse. Search date: October 1, 2015.

The Authors

MARC A. CHILDRESS, MD, is the associate director of the Primary Care Sports Medicine Fellowship at Virginia Commonwealth University Fairfax Family Practice.

BLAIR A. BECKER, MD, is a faculty physician at the Group Health Family Medicine Residency, Seattle, Wash.

Address correspondence to Marc A. Childress, MD, Fairfax Family Practice, 3650 Joseph Siewick Dr., Ste. 400, Fairfax, VA 22033 (e-mail: mchildress@ffpcs.com). Reprints are not available from the authors.

REFERENCES

- Bono CM, Ghiselli G, Gilbert TJ, et al.; North American Spine Society. An evidence-based clinical guideline for the diagnosis and treatment of cervical radiculopathy from degenerative disorders. *Spine J*. 2011;11(1):64-72.
- Radhakrishnan K, Litchy WJ, O'Fallon WM, Kurland LT. Epidemiology of cervical radiculopathy. A population-based study from Rochester, Minnesota, 1976 through 1990. *Brain*. 1994;117(pt 2):325-335.
- Levine MJ, Albert TJ, Smith MD. Cervical radiculopathy: diagnosis and nonoperative management. *J Am Acad Orthop Surg*. 1996;4(6):305-316.
- Rothman RH, Marvel JP Jr. The acute cervical disk. *Clin Orthop Relat Res*. 1975;(109):59-68.
- Kuijper B, Tans JT, van der Kallen BF, Nollet F, Lycklama A Nijeholt GJ, de Visser M. Root compression on MRI compared with clinical findings in patients with recent onset cervical radiculopathy. *J Neurol Neurosurg Psychiatry*. 2011;82(5):561-563.
- Ghasemi M, Golabchi K, Mousavi SA, et al. The value of provocative tests in diagnosis of cervical radiculopathy. *J Res Med Sci*. 2013;18(suppl 1):S35-S38.
- Vallée JN, Feydy A, Carlier RY, Mutschler C, Mompont D, Vallée CA. Chronic cervical radiculopathy: lateral-approach periradicular corticosteroid injection. *Radiology*. 2001;218(3):886-892.
- Eubanks JD. Cervical radiculopathy: nonoperative management of neck pain and radicular symptoms. *Am Fam Physician*. 2010;81(1):33-40.

Cervical Radiculopathy

9. Rubinstein SM, Pool JJ, van Tulder MW, Riphagen II, de Vet HC. A systematic review of the diagnostic accuracy of provocative tests of the neck for diagnosing cervical radiculopathy. *Eur Spine J*. 2007;16(3):307-319.
10. Shah KC, Rajshekhar V. Reliability of diagnosis of soft cervical disc prolapse using Spurling's test. *Br J Neurosurg*. 2004;18(5):480-483.
11. Wainner RS, Fritz JM, Irrgang JJ, Boninger ML, Delitto A, Allison S. Reliability and diagnostic accuracy of the clinical examination and patient self-report measures for cervical radiculopathy. *Spine (Phila Pa 1976)*. 2003;28(1):52-62.
12. Shabat S, Leitner Y, David R, Folman Y. The correlation between Spurling test and imaging studies in detecting cervical radiculopathy. *J Neuroimaging*. 2012;22(4):375-378.
13. Friedenbergs ZB, Miller WT. Degenerative disk disease of the cervical spine. *J Bone Joint Surg Am*. 1963;45:1171-1178.
14. Modic MT, Masaryk TJ, Ross JS, Mulopulos GP, Bundschuh CV, Bohlman H. Cervical radiculopathy: value of oblique MR imaging. *Radiology*. 1987;163(1):227-231.
15. Spurling RG, Segerberg LH. Lateral intervertebral disk lesions in the lower cervical region. *J Am Med Assoc*. 1953;151(5):354-359.
16. Gross A, Kay TM, Paquin JP, et al.; Cervical Overview Group. Exercises for mechanical neck disorders. *Cochrane Database Syst Rev*. 2015;(1):CD004250.
17. Boyles R, Toy P, Mellon J Jr, Hayes M, Hammer B. Effectiveness of manual physical therapy in the treatment of cervical radiculopathy: a systematic review. *J Man Manip Ther*. 2011;19(3):135-142.
18. Kuijper B, Tans JT, Beelen A, Nollet F, de Visser M. Cervical collar or physiotherapy versus wait and see policy for recent onset cervical radiculopathy: randomised trial. *BMJ*. 2009;339:b3883.
19. Teresi LM, Lufkin RB, Reicher MA, et al. Asymptomatic degenerative disk disease and spondylosis of the cervical spine: MR imaging. *Radiology*. 1987;164(1):83-88.
20. Bartlett RJ, Hill CR, Gardiner E. A comparison of T2 and gadolinium enhanced MRI with CT myelography in cervical radiculopathy. *Br J Radiol*. 1998;71(841):11-19.
21. Cleland JA, Fritz JM, Whitman JM, Heath R. Predictors of short-term outcome in people with a clinical diagnosis of cervical radiculopathy. *Phys Ther*. 2007;87(12):1619-1632.
22. Persson LC, Carlsson CA, Carlsson JY. Long-lasting cervical radicular pain managed with surgery, physiotherapy, or a cervical collar. A prospective, randomized study. *Spine (Phila Pa 1976)*. 1997;22(7):751-758.
23. Thoomes EJ, Scholten-Peeters W, Koes B, Falla D, Verhagen AP. The effectiveness of conservative treatment for patients with cervical radiculopathy: a systematic review. *Clin J Pain*. 2013;29(12):1073-1086.
24. Graham N, Gross A, Goldsmith CH, et al. Mechanical traction for neck pain with or without radiculopathy. *Cochrane Database Syst Rev*. 2008;(3):CD006408.
25. van der Heijden GJ, Beurskens AJ, Koes BW, Assendelft WJ, de Vet HC, Bouter LM. The efficacy of traction for back and neck pain: a systematic, blinded review of randomized clinical trial methods. *Phys Ther*. 1995;75(2):93-104.
26. Young IA, Michener LA, Cleland JA, Aguilera AJ, Snyder AR. Manual therapy, exercise, and traction for patients with cervical radiculopathy: a randomized clinical trial [published corrections appear in *Phys Ther*. 2009;89(11):1254-1255, and *Phys Ther*. 2010;90(5):825]. *Phys Ther*. 2009;89(7):632-642.
27. Pain in the neck and arm: a multicenter trial of the effects of physiotherapy, arranged by the British Association of Physical Medicine. *Br Med J*. 1966;1(5482):253-258.
28. Klaber Moffett JA, Hughes GI, Griffiths P. An investigation into the effects of cervical traction. Part 1: clinical effectiveness. *Clin Rehabil*. 1990;4:205-211.
29. Fritz JM, Thackeray A, Brennan GP, Childs JD. Exercise only, exercise with mechanical traction, or exercise with over-door traction for patients with cervical radiculopathy, with or without consideration of status on a previously described subgrouping rule: a randomized clinical trial. *J Orthop Sports Phys Ther*. 2014;44(2):45-57.
30. Jellad A, Ben Salah Z, Boudokhane S, Migaou H, Bahri I, Rejeb N. The value of intermittent cervical traction in recent cervical radiculopathy. *Ann Phys Rehabil Med*. 2009;52(9):638-652.
31. Swezey RL, Swezey AM, Warner K. Efficacy of home cervical traction therapy. *Am J Phys Med Rehabil*. 1999;78(1):30-32.
32. Patel KC, Gross A, Graham N, et al. Massage for mechanical neck disorders. *Cochrane Database Syst Rev*. 2012;(9):CD004871.
33. Peloso P, Gross A, Haines T, et al.; Cervical Overview Group. Medicinal and injection therapies for mechanical neck disorders. *Cochrane Database Syst Rev*. 2007;(3):CD0000319.
34. Ghasemi M, Maseali A, Rezvani M, Shaygannejad V, Golabchi K, Norouzi R. Oral prednisolone in the treatment of cervical radiculopathy: a randomized placebo controlled trial. *J Res Med Sci*. 2013;18(suppl 1):S43-S46.
35. Kwon JW, Lee JW, Kim SH, et al. Cervical interlaminar epidural steroid injection for neck pain and cervical radiculopathy: effect and prognostic factors. *Skeletal Radiol*. 2007;36(5):431-436.
36. Diwan S, Manchikanti L, Benyamin RM, et al. Effectiveness of cervical epidural injections in the management of chronic neck and upper extremity pain. *Pain Physician*. 2012;15(4):E405-E434.
37. Stav A, Ovadia L, Sternberg A, Kaadan M, Weksler N. Cervical epidural steroid injection for cervicobrachialgia. *Acta Anaesthesiol Scand*. 1993;37(6):562-566.
38. Bush K, Hillier S. Outcome of cervical radiculopathy treated with periradicular/epidural corticosteroid injections: a prospective study with independent clinical review. *Eur Spine J*. 1996;5(5):319-325.
39. Castagnera L, Maurette P, Pointillart V, Vital JM, Erny P, S negas J. Long-term results of cervical epidural steroid injection with and without morphine in chronic cervical radicular pain. *Pain*. 1994;58(2):239-243.
40. Cicala RS, Thoni K, Angel JJ. Long-term results of cervical epidural steroid injections. *Clin J Pain*. 1989;5(2):143-145.
41. Honet JC, Puri K. Cervical radiculitis: treatment and results in 82 patients. *Arch Phys Med Rehabil*. 1976;57(1):12-16.
42. R s nen P, Ohman J, Sintonen H, et al. Cost-utility analysis of routine neurosurgical spinal surgery. *J Neurosurg Spine*. 2006;5(3):204-209.