



**Key:**  
**C** = Compatible  
**I** = Incompatible  
**Cs** = Compatible Only if Drugs mixed in NS  
**CD** = Compatible Only if Drugs mixed in Dextrose  
**C\*** = Conc./Time dependent,  
**CALL PHARMACY**

	dextrose IV soln	saline IV Soln	dex/Na/K IV soln	< 1 unit/mL heparin	PN (NO heparin)	PN (with heparin)	intraipids (fats)	alprostadil (PGE1)	amiodarone	calcium CHLORIDE	calcium GLUCONATE	cisatracurium	cycloSPORINE	dexmedetomidine	DOBUTamine	DOPamine	EPINEPHrine	esmolol	fentanyl	heparin > 1 unit/mL	HYDROMorphone	insulin	isoproterenol	ketamine	KCl (Potassium)	labetalol	lidocaine	methy/PREDNISolone	midazolam	milrinone	morphine	naloxone	nICARdipine	nitroGLYCERIN	nitroPRUSSIDE	norepinephrine	phenylephrine	remifentanil	sodium bicarb.	TACROlimus	terbutaline	tobramycin	vasopressin	vecuronium						
intralipid (Fats)	C	C	C	C	C	C	C	C	C	I	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	I	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C						
isoproterenol	C	C	C	C	C	C	C	C	C*	C	C	C	C	C	C	C	C	C	C	C	C	I	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C					
KCl (potassium)	C	C	C	C	C	C	C	C	C*	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C					
ketamine	C	C	C	I					C*	C	C	C	C	C	C	C	C	C	C	C	C	I	C	I	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C					
ketorolac	C	C	C	C					C*	I	C	C	C	C	I	C	C	C	I	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C					
labetalol	C	C	C						C*	C	C	C	C	C	C	C	C	C	C	C	C*	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C					
lidocaine	C	C	C	C	C	C	C	C	C*	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C				
lorazepam	C	C	C	C	C	C	C		C*	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C				
magnesium sulf.	C	C	C	C	C	C	C			I	I	C	CS	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C				
mannitol	C	C	C	C	C	C	C			C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C				
meropenem	C	C	C	C	C	C	C			C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C				
methyIPREDNISolone	C	C	C	C	C	C	C	C	C*	I	I	C	C	C	C	C	C	CS	C	C	C	C	C	C*	C	C	C	C	C	C	C	C	C	C	C	C	CD	C	C	C	C	C	C	C	C	C				
metoclopramide	C	C	C	C	C	C	C	C	C	I	I	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C				
metRONIDAZOLE	C	C	C	C	C	C	C		C	I	C	C	C	C	C	CS	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C			
midazolam	C	C	C	C	C*	C*	I	C	C*	C	C	C	C	C	C	C	C	C	C	C	C	C	CD	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C			
micafungin	C	C	C	C	C	C	C			I	C	C	I	C		I	C	I	C		C	C	I			C	I	C		I	C	C	C	C	C							C	C	C	C	C	C			
milrinone	C	C	C	C	C	C	C		C*	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C			
morphine	C	C	C	C	C	C	C	C	C*	C	C	C	C	C	C	C	C	C	C	C	C	C	I	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C			
mycophenolate	C			I					CD	I		C*	C	I	CD	CD	CD	CD	CD	I	CD	CD	CD			CD	CD	CD	I	CD	CD	CD	CD	CD	I	CD	CD	I	CD	CD	I	CD	CD	I	CD	CD	I	CD		
naloxone	C	C	C	C	C	C				C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C		
NICARdipine	C	C	C							C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C		
nitroGLYCERIN	C	C	C	C	C	C	C	C	C*	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C		
nitroPRUSSIDE	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	CD	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	
norepinephrine	C	C	C	C	C	C	C		C*	C	C	C	C	C	C	C	C	C	C	C	C	I	C			C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C		
octreotide	C	C	C	C	C	C	I		C*	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C		
ondansetron	C	C	C	C	C	C	I			C	C	C	C	C	C	C	C	C	C	C	C	CD				C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C		
oxacillin	C	C	C	C	C	C	C			I	I		C		I	C	C	I	C	C	C	C	C	C			C	C	C		C	C	C	C	C	C	C	C	CD	C	C	C	C	C	I	C				
pancuronium	C	C	C	C	C	C	C			C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C		
pantoprazole	C	C	C		I	I			C	I	I	I	I	I	I	I	I	I	I	I	I	I	I	I	I	I	I	I	I	I	I	I	I	I	I	I	I	I	I	I	I	I	I	I	I	I	I	I	I	
penicillin G K	C	C	C	C	C	C	C	C	C*	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C		
PENTObarbital	C	C	C	C	I	I	I			C	C		I	C	I	C	I	I	C	C	C	C	I			C	C	I	C	I	C	I	C	I	C	I	C	C	I	C	I	C	I	C	I	C	I	C	I	CD
PHENObarbital	C	C	C	C	I	I	I			C	C		I	C	I	C	I	I	C	C	C	C	C	I			C	C	I	C	I	C	I	C	I	C	C	I	C	I	C	I	C	I	C	I	C	I	C	
phenylephrine	C	C	C	C	C	C	C		C*	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	
phosphate(Na or K)	C	C	C	C	I	I	I			I	I	I		C	I	C		C		C	C	C			I	C	C																							
piperacillin / zosyn	C	C	C	C	C	C	C			I	C	C	C	C	I	C	C	C	C	C	C	I	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C		
PN (with OR without Heparin)	C	C	C	C	C	C	C	C		C	I	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C		
propranolol	C	C	C	C						C	C	C	C	C	C	C	C	C	C	C	C	I	C			C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C		
ranitidine	C	C	C	C	C	C	C	C		C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	
remifentanil	C	C	C	C						C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	
rocuronium	C	C	C	C			I		C*	C	C	C	C	C	C	C	C	C	C	C	C	I	C			C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C*		
sodium bicarb.	C	C	C	C	I	I	C			I	I	I	I	C	C	I	CD	I	C	C	C	I	C	I	I	C	I	C	C	I	C	C	I	C	C	I	C	C	I	C	C	C	C	C	C	C	C	C	C	
TACROlimus	C	C	C	C	C	C	C			C*	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	
terbutaline	C	C	C	C	C	C	C			C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	
tobramycin	C	C	C	C	C	C	C			C*	C	C	C	C	C	C	C	C	C	C	C	I	C	I	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	
vancomycin	C	C	C	C	C	C	C	C*	C*	C	C	C	C	C	C	C	C	C	C	C	C	I	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C		
vasopressin	C	C	C	C	C	C	C			C*	C	C	C	C	C	C	C	C	C	C	C	I	C			C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	
vecuronium	C	C	C	C	C	C	C	C		C*	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	
voriconazole	C	C	C	C						C*	C																																							

Additional information for RX use only. Not available on nursing charts.

Key:	dextrose IV soln	saline IV Soln	dex/Na/K IV soln	< 1 unit/mL heparin	PN (NO heparin)	PN (with heparin)	intraIipids (fats)	alprostadil (PGE1)	amiodarone	calcium CHLORIDE	calcium GLUCONATE	cisatracurium	cycloSPORINE	dexmedetomidine	DOBUTamine	DOPamine	EPINEPHrine	esmolol	fentanyl	heparin > 1 unit/mL	HYDROmorphone	insulin	isoproterenol	KCl (Potassium)	ketamine	labetalol	lidocaine	methyIPREDNISolone	midazolam	milrinone	morphine	naloxone	niCARdipine	nitroGLYCERIN	nitroPRUSSIDE	norepinephrine	phenylephrine	remifentanyl	sodium bicarb.	TACROLimus	terbutaline	tobramycin	vasopressin	vecuronium						
acetaZOLAMIDE	C	C	C		I	I	I																																											
albumin	C	C	C				I																		C				I																					
alteplase	I	C	I												I	I				I								C																						
ambisome	C	I	I	CD						I	I		I	I	I	I	CD	CD	CD	CD	C		CD	CD		I	CD	CD	I	CD	I	I	I	I	CD	CD		CD	I	I	CD		I	CD		I	CD	I		
argatroban	C	C	C	C					I	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C		
bivalirudin	C	C	C	C					I	C	C	C	C	C	C*	C	C	C	C	C	C	C	C	C		C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C		
butorphanol										C	C	C	C	C	C	C	C	C	C	C		I	C	C		C	C	C	C*	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C		
cefurOXIME	C	C	C	C	C	C		C	C*	I	C	I	C	C	I	C	C	C	C	C	C	C	C	C	C	I	C	C	I	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C		
chlorothiazide	C	C	C		I	I																					C																							
doxapram	C	C	C	C						C	C				I	C	C		C									I																				C	C	
foscarnet	C	C	C	C	C	C			I	I	I	C	C	C	I	C	C	C	C	C	C	C	C	C		I	C	I	I	C	C	C	I	C	C	I	C	C	C	C	C	C	C	C	C	C	C	C	C	
fosphenytoin	C	C	C											C										C					I																					
ibuprofen lysine	C	C	C	C	I	I									I	I	C			C		C	I	C					I		C*																		I	
levetiracetam	C	C	C																																															
methadone		C																																																
neostigmine	C	C	C	C																			C		C																									
nesiritide	C	C	C	I					C*	C	C	C	C	C				C	C	I	C	I		C		I		C	C	C		C	C	C	C	C			C	C	C		C	C	C		C	C		
procainamide	I	C	I	CS					C*	CS	CS	C	CS	CS	C	C	C	C	C	C	C	C	C	C		C	C	C	C	C	I	C																	C	
propofol	C		C	CD	C	C				I	CD		CD	C	CD	CD	CD	CD	CD	CD	CD	CD	CD	CD	C	CD		I		CD																				
quiNIDine gluconate	C			C*					I	CD	C		C	C	C	C	C	C	C	C*	C	I	C		C	C	I	CD	C	C	C																			
urokinase	C	C	C		C	C				C	C		C	C	C	C	C	C	C	C	C	C	C		C	C	C	C	C	C	C																			

© 2014 Bloomberg Children's Center Pharmacy, Guidelines for Y-Site use only; Authors not responsible for continued currency of information, errors, or omissions.