



Algebra 1

Transitional Curriculum REVISED 2012

BLACKLINE MASTERS

LOUISIANA DEPARTMENT OF EDUCATION

Unit 1, Activity 1, Identifying and Classifying Numbers

Identifying and Classifying Numbers

1. Explain the difference between a rational and an irrational number.

Classify the following numbers as rational or irrational.

4. $\sqrt{6}$ 5. $\sqrt{16}$ 2. $\frac{1}{2}$ 3.8 6. *π*

7. List the set of all natural numbers.

8. List the set of whole numbers less than 4.

9. List the set of integers such that -3 < x < 5.

Classify the following numbers as rational, irrational, natural, whole and/or integer. (A number may belong to more than one set)

12. $4\frac{2}{3}$ 13. $\sqrt{3}$ 10. -3 14. 0

15. Using the following set of numbers: $A = \{\sqrt{3.6}, 0.36, -\frac{3}{6}, 0.3\overline{6}, 0, 3^6, -3, \sqrt{36}, 3.63363336...\}$, place each element in the appropriate subset. (Numbers may belong to more than one subset)

rational numbers_____

natural numbers_____

irrational numbers_____

integers_____

True or False?

16. All whole numbers are rational numbers.

17. All integers are irrational numbers.

18. All natural numbers are integers.

whole numbers_____

Unit 1, Activity 1, Identifying and Classifying Numbers with Answers

Identifying and Classifying Numbers

1. Explain the difference between a rational and an irrational number. A rational number can be expressed as the ratio of two integers. An irrational number is any real number that is not rational

Classify the following numbers as rational or irrational.

- 4. $\sqrt{6}$ 5. $\sqrt{16}$ 6. π *irrational rational irr* 2. $\frac{1}{2}$ 3.8 rational irrational rational 7. List the set of all natural numbers. $\{1, 2, 3...\}$
- 8. List the set of whole numbers less than 4. $\{0, 1, 2, 3\}$
- 9. List the set of integers such that -3 < x < 5. $\{-2, -1, 0, 1, 2, 3, 4\}$

Classify the following numbers as rational, irrational, natural, whole and/or integer. (A number may belong to more than one set)

10. -3 rational 12. $4\frac{2}{3}$ rational 13. $\sqrt{3}$ irrational 14. 0 rational, integer whole number integer 15. Using the following set of numbers:

 $A = \{\sqrt{3.6}, 0.36, -\frac{3}{6}, 0.3\overline{6}, 0, 3^6, -3, \sqrt{36}, 3.63363336...\}$, place each element in the appropriate subset. (Numbers may belong to more than one subset)

rational numbers $\{\sqrt{3.6}, 0.36, -\frac{3}{6}, 0.3\overline{6}, 0, 3^6, -3, \sqrt{36}\}$ irrational numbers <u>3.63363336</u>


natural numbers $\sqrt{36}, 3^6$ whole numbers $0, \sqrt{36}, 3^6$

integers___-3, 0, $\sqrt{36}$, 3^6 _____

True or False?

16. All whole numbers are rational numbers. True

- 17. All integers are irrational numbers. False.
- 18. All natural numbers are integers. *True*



Lamp Flowchart

What is a Flowchart?

Flowchart Definitions and Objectives:

Flowcharts are maps or graphical representations of a process. Steps in a process are shown with symbolic shapes, and the flow of the process is indicated with arrows connecting the symbols. Computer programmers popularized flowcharts in the 1960s, using them to map the logic of programs. In quality improvement work, flowcharts are particularly useful for displaying how a process currently functions or could ideally function. Flowcharts can help you see whether the steps of a process are logical, uncover problems or miscommunications, define the boundaries of a process, and develop a common base of knowledge about a process. Flowcharting a process often brings to light redundancies, delays, dead ends, and indirect paths that would otherwise remain unnoticed or ignored. But flowcharts don't work if they aren't accurate.

A flowchart (also spelled flow-chart and flow chart) is a schematic representation of a process. It is commonly used in business/economic presentations to help the audience visualize the content better, or to find flaws in the process.

The flowchart is one of the seven basic tools of quality control, which include the histogram, Pareto chart, check sheet, control chart, cause-and-effect diagram, flowchart, and scatter diagram. Examples include instructions for a bicycle's assembly, an attorney who is outlining a case's timeline, diagram of an automobile plant's work flow, or the decisions to make on a tax form.

Generally the start point, end points, inputs, outputs, possible paths and the decisions that lead to these possible paths are included.

Flow-charts can be created by hand or manually in most office software, but lately specialized diagram drawing software has emerged that can also be used for the intended purpose. See below for examples.

Flowchart History:

Flowcharts were used historically in electronic data processing to represent the conditional logic of computer programs. With the emergence of structured programming and structured design in the 1980s, visual formalisms like data flow diagrams and structure charts began to supplant the use of flowcharts in database programming. With the widespread adoption of such ALGOL-like computer languages as Pascal, textual models have been used more and more often to represent algorithms. In the 1990s Unified Modeling Language began to synthesize and codify these modeling techniques.

Today, flowcharts are one of the main tools of business analysts and others who seek to describe the logic of a process in a graphical format. Flowcharts and cross-functional flowcharts can commonly be found as a key part of project documentation or as a part of a

Unit 1, Activity 2, What is a Flowchart?

business process document. Flowcharts are widely used in education, clinical settings, service industries and other areas where graphical, logical depiction of a process is helpful.

When should flowcharts be used?

At the beginning of your process improvement efforts, an "as-is" flowchart helps your team and others involved in the process to understand how it currently works. The team may find it helpful to compare this "as-is flowchart" with a diagram of the way the process is supposed to work. Later, the team will develop a flowchart of the modified process again, to record how it actually functions. At some point, your team may want to create an ideal flowchart to show how you would ultimately like the process to be performed.

Among the benefits of using flowcharts are that they:

- **Promote process understanding by explaining the steps pictorially.** People may have differing ideas about how a process works. A flowchart can help you gain agreement about the sequence of steps. Flowcharts promote understanding in a way that written procedures cannot. One good flowchart can replace pages of words.
- **Provide a tool for training employees.** Because of the way they visually lay out the sequence of steps in a process, flowcharts can be very helpful in training employees to perform the process according to standardized procedures.
- Identify problem areas and opportunities for process improvement. Once you break down the process steps and diagram them, problem areas become more visible. It is easy to spot opportunities for simplifying and refining your process by analyzing decision points, redundant steps, and rework loops.
- **Depict customer-supplier relationship,** helping the process workers understand who their customers are, and how they may sometimes act as suppliers, and sometimes as customers in relation to other people.

What symbols are used in flowcharts?

The symbols that are commonly used in flowcharts have specific meanings and are connected by arrows indicating the flow from one step to another.

- Oval. An oval indicates both the starting point and the ending point of the process.
- Box. A box represents an individual step or activity in the process.
- **Diamond.** A diamond shows a decision point, such as yes/no or go/no-go. Each path emerging from the diamond must be labeled with one of the possible answers.
- **Circle.** A circle indicates that a particular step is connected within the page. A numerical value is placed in the circle to indicate the sequence continuation.
- **Pentagon.** A pentagon indicates that a particular step of the process is connected to another page or part of the flowchart. A letter placed in the circle clarifies the continuation.
- Flow line. This indicates the direction flow of the process.

Excerpt taken from: http://www.edrawsoft.com/Flowchart-tutorial.php

Unit 1, Activity 2, DL-TA

DL-TA for (title)_____

Prediction question(s):_____

Using the title, your own background knowledge, and any other contextual clues, make your predictions.


Before reading:

During reading:

During reading:

During reading:

After reading:



Unit 1, Activity 4, Classifying Numbers: Sums and Products of Rational and Irrational Numbers

Number	Irrational	Rational	Integer	Whole	Natural
Example: 2		X	X	X	X
A6					
B. $\frac{3\pi}{4}$					
C. 17.3					
D. 3√2					
E. √16					
F. 8					
G. $\frac{3}{4}$					
H. 5.0					
I17.3					
J. 2√2					

Part I: Classify the real numbers below in all possible ways.

Part II. Perform the indicated operations. Classify each sum or product as rational or irrational by circling the appropriate classification.

Example: A + C	-6 + 17.3 = 11.3	rational	irrational
1) B x G		rational	irrational
2) G + A		rational	irrational
3) E x C		rational	irrational
4) I x D		rational	irrational
5) G + B		rational	irrational
6) E x F		rational	irrational
7) I + J		rational	irrational
8) E + A		rational	irrational
9) F + J		rational	irrational
10) G x A		rational	irrational

Unit 1, Activity 4, Classifying Numbers: Sums and Products of Rational and Irrational Numbers

Part III: Using your answers from Part II, answer the following questions with complete sentences. Justify your classification using appropriate algebraic language.

A. What is the classification of the sum of two rational numbers?

B. What is the classification of the product of two rational numbers?

C. What is the classification of the sum of a rational and an irrational number?

D. What is the classification of the product of a rational and an irrational number?

E. When is it possible for the product of a rational and an irrational number to result in a rational value?

Unit 1, Activity 4, Classifying Numbers: Sums and Products of Rational and Irrational Numbers with Answers

Number	Irrational	Rational	Integer	Whole	Natural
Example: 2		X	X	X	X
A6		X	X		
B. $\frac{3\pi}{4}$	X				
C. 17.3		X			
D. 3√2	X				
E. $\sqrt{16}$		X	X	X	X
F. 8		X	X	X	X
G. $\frac{3}{4}$		X			
H. 5.0		X	X	X	X
I17.3		X			
J. 2√2	X				

Part I: Classify the real numbers below in all possible ways.

Part II. Perform the indicated operations. Classify each sum or product as rational or irrational by circling the appropriate classification.

Example: A + C	-6 + 17.3 = 11.3	rational	irrational
1) B x G	$\frac{3\pi}{4} \times \frac{3}{4} = \frac{9\pi}{16}$	rational	irrational
2) G + A	$\frac{3}{4} + (-6) = -5\frac{1}{4}$	rational	irrational
3) E x C	$\sqrt{16} \times 17.3 = 69.2$	rational	irrational
4) I x D	$-17.3 \times 3\sqrt{2} = -51.9\sqrt{2}$	rational	irrational
5) G + B	$\frac{3}{4} + \frac{3\pi}{4} = \frac{3+3\pi}{4}$	rational	irrational
6) E x F	$\sqrt{16} \times 8 = 32$	rational	irrational
7) I + J	$-17.3 + 2\sqrt{2}$	rational	irrational
8) E + A	$\sqrt{16} + -6 = -2$	rational	irrational
9) F + J	$8 + 2\sqrt{2}$	rational	irrational
10) G x A	$\frac{3}{4} \times -6 = -\frac{9}{2}$	rational	irrational

Unit 1, Activity 4, Classifying Numbers: Sums and Products of Rational and Irrational Numbers with Answers

Part III: Using your answers from Part II, answer the following questions with complete sentences. Justify your classification using appropriate algebraic language.

A. What is the classification of the sum of two rational numbers? *The sum of two rational numbers is rational because the sum can be written as a fraction.*

B. What is the classification of the product of two rational numbers? *The product of two rational numbers is a rational number because the product can be written as a fraction.*

C. What is the classification of the sum of a rational and an irrational number? *The sum of a rational and an irrational number is irrational since the sum cannot be written as a fraction in which the numerator and denominator are both integers.*

D. What is the classification of the product of a rational and an irrational number? *The product of a rational and an irrational number is irrational because the product cannot be written as a fraction in which the numerator and denominator are both integers.*

E.. When is it possible for the product of a rational and an irrational number to result in a rational value?

The product of rational and an irrational number can result in a rational value when one of the factors is the rational number 0.

Unit 1, Activity 4, Classifying Numbers: Sums and Products of Rational and Irrational Numbers Homework

Homework Assignment: Classification of Sums and Products

1. Find each sum or product. Determine its classification as rational or irrational.

a.	1000 + (-288) = _	
b.	54.25 + $\sqrt{26}$ =	
c.	0.7 × 17 =	
d.	4.1 x √49 =	
e.	-36 x √3 =	

2. Write an example for each condition.a. The product of two rational numbers:

.

- b. The sum of two rational numbers:
- c. The product of a rational and an irrational number:
- d. The sum of a rational and an irrational number.

Unit 1, Activity 4, Classifying Numbers: Sums and Products of Rational and Irrational Numbers Homework with Answers

1. Find each sum or product. Determine its classification as rational or irrational.

a. $1000 + (-288) = 712 \ rational$ b. $54.25 + \sqrt{26} = 54.25 + \sqrt{26} \ irrational$ c. $0.7 \times 17 = 11.9 \ rational$ d. $4.1 \ge \sqrt{49} = 28.7 \ rational$

e. -36 x $\sqrt{3} = \frac{-36}{\sqrt{3}} \sqrt{3}$ irrational

Write an example for each condition.
 a. The product of two rational numbers:

Answers will vary.

b. The sum of two rational numbers:

Answers will vary.

c. The product of a rational and an irrational number:

Answers will vary.

d. The sum of a rational and an irrational number.

Answers will vary.

An Activity Comparing Foot Length and Shoe Size

Student Worksheet

Using the standard measuring edge of your ruler, measure and record the length of the student's foot, in centimeters, and the student's shoe size. Record 10 sets of data for girls and 10 sets for boys.

Length of Foot	Shoe Size	Ratio of Foot Length to Shoe Size	Length of Foot	Shoe Size	Ratio of Foot Length to Shoe Size

GIRLS	
-------	--

BOYS

Answer these questions:

- 1. Which is the independent and which is the dependent variable? How do you know?
- 2. Write ordered pairs on the line below, graph them, and look for a pattern.

- 3. Does the data appear to be linear? Explain your reasoning.
- 4. Using your calculator, find the average ratio of foot length to shoe size. This is the constant of variation. What does this value mean in the context of the data you have collected?

- 5. Write an equation that models the situation. _____
- 6. List three other examples of positive relationships from real-life.

Unit 1, Activity 5, Foot Length and Shoe Size

An Activity to Study Dimension of a Rectangle

Student Worksheet

Each group has 36 algebra unit tiles. Using all of the tiles you have been given, arrange the tiles in a rectangle and record the height and width. Continue to make rectangles until you think you have created all possible rectangles. Record the area for each rectangle.

Height (h)	Width (w)	Area

Answer these questions:

1. Does it matter which variable is independent and which variable is dependent? Explain.

2. Get together with other groups and decide which the entire class will use. Record the class decision below.

3. Using the table and the class decision about independent and dependent variables, write the ordered pairs below.

4. Graph these pairs on the paper provided and look for relationships in the graphed data.

5. Write an equation to model this situation.


Unit 1, Activity 6, Exponential Growth and Decay

Exponential Growth and Decay

In this activity, fold a piece of computer paper in half as many times as possible. After each fold, stop to fill in a row of the table. Continue folding and recording until the table is filled.

NUMBER OF FOLDS	NUMBER OF REGIONS	AREA OF SMALLEST REGION
0		
1		
2		
3		
4		
5		
6		
7		
N		

Part I: Label the axes of the graph below before you plot your data. One axis is for Number of Folds and one axis is for Number of Regions. Think about the scale of your range and plan your markings before you begin.




Unit 1, Activity 6, Exponential Growth and Decay

	swer the following questions with the class: Identify the independent and the dependent variables. The independent variable is	
	The dependent variable is	
2.	Is the graph linear? Explain your reasoning.	
3.	Describe the pattern that occurs in the data.	

4. Write an equation to determine the number of regions if there are n folds.

Part II: Label the axes of the graph below before you plot your data. One axis is for Number of Folds and one axis is Area of Smallest Region. Think about the scale of your range and plan your markings before you begin.



Unit 1, Activity 6, Exponential Growth and Decay

Answer the following questions with the class:

- Identify the independent and the dependent variables. The independent variable is ______
 The dependent variable is ______
- 2. Is the graph linear? Explain your reasoning.

3. Describe the pattern that occurs in the data.

4. Write an equation to determine the number of regions if there are *n* folds.


Unit 1, Activity 6, Exponential Growth and Decay with Answers

Exponential Growth and Decay

In this activity, fold a piece of computer paper in half as many times as possible. After each fold, stop to fill in a row of the table. Continue folding and recording until the table is filled

NUMBER OF FOLDS	NUMBER OF REGIONS	AREA OF SMALLEST REGION
0	1	1
1	2	$\frac{1}{2}$ or 2^{-1}
2	4	$\frac{1}{4}$ or 2^{-2}
3	8	$\frac{1}{8}$ or 2^{-3}
4	16	$\frac{1}{16}$ or 2^{-4}
5	32	$\frac{1}{32}$ or 2^{-5}
6	64	$\frac{1}{64}$ or 2^{-6}
7	128	$\frac{1}{128}$ or 2^{-7}
N	2^n	$\frac{1}{2^n}$ or 2^{-n}

Part I: Label the axes of the graph below before you plot your data. One axis is for Number of Folds and one axis is for Number of Regions. Think about the scale of your range and plan your markings before you begin.



Unit 1, Activity 6, Exponential Growth and Decay with Answers

Answer the following questions with the class:

 1. Identify the independent and the dependent variables.

 The independent variable is
 number of folds

 The dependent variable is
 number of regions


2. Is the graph linear? Explain your reasoning. *The graph is not linear because the data points do not seem to form a line; there is not a constant rate of change.*

3. Describe the pattern that occurs in the data. <u>*The number of regions increases by a power of two. The power is equal to the number of folds.*</u>

4. Write an equation to determine the number of regions if there are *n* folds.

 $r = 2^{n}$

Part II: Label the axes of the graph below before you plot your data. One axis is for Number of Folds and one axis is Area of Smallest Region. Think about the scale of your range and plan your markings before you begin.



Answer the following questions with the class:

 1. Identify the independent and the dependent variables.

 The independent variable is
 number of folds

 The dependent variable is
 area of smallest region

Unit 1, Activity 6, Exponential Growth and Decay with Answers

2. Is the graph linear? Explain your reasoning. *The graph is not linear because the data points do not seem to form a line; there is not a constant rate of change.*

3. Describe the pattern that occurs in the data. <u>The area of the smallest region decreases by a</u> <u>negative power of two (or a power of $\frac{1}{2}$). The power is equal to the number of folds.</u>

4. Write an equation to determine the number of regions if there are *n* folds. $r = 2^{-n}$

Pay Day!

In your math log, respond to the following question.

Part I:

Which of the following jobs would you choose? Give reasons to support your answer.

- Job A: Salary of \$1 for the first year, \$2 for the second year, \$4 for the third year, continuing for 25 years.
- Job B: Salary of \$1 Million a year for 25 years

Part II:

At the end of 25 years, which job would produce the largest amount in total salary?

NUMBER OF	YEARLY	TOTAL	YEARLY	TOTAL
YEARS WORKED	SALARY	SALARY	SALARY	SALARY
	JOB A	JOB A	JOB B	JOB B
1	\$1	\$1	\$1,000,000	\$1,000,000
2				
2				
3				
4				
5				
6				
N				

- 1. Predict when the salaries will be equal. Explain your reasoning.
- 2. Does Job A represent linear or exponential growth? Explain your reasoning.
- 3. Does Job B represent linear or exponential growth? Explain your reasoning.
- 4. Write an equation to represent the salary for each job option.

Pay Day!

In your math log, respond to the following question.

Which of the following jobs would you choose?

Job A: Salary of \$1 for the first year, \$2 for the second year, \$4 for the third year, continuing for 25 years.

Job B: Salary of \$1 Million a year for 25 years

Give reasons to support your answer.

At the end of 25 years, which job would produce the largest amount in total salary?

NUMBER OF	YEARLY	TOTAL	YEARLY	TOTAL
YEARS WORKED	SALARY	SALARY	SALARY	SALARY
	JOB A	JOB A	JOB B	JOB B
1	\$1	\$1		
1	ΦI	Φ1	\$1,000,000	\$1,000,000
2	\$2	\$3	\$1,000,000	\$2,000,000
Z	\mathcal{P}	φς	\$1,000,000	\$2,000,000
2	¢ 4	ф 7	¢1,000,000	¢2,000,000
3	\$4	\$7	\$1,000,000	\$3,000,000
4	¢o	¢15	¢1,000,000	\$1,000,000
4	\$8	\$15	\$1,000,000	\$4,000,000
	¢16	¢21	¢1,000,000	<i>#<i>F</i> 000 000</i>
5	\$16	\$31	\$1,000,000	\$5,000,000
	¢ 22	¢(2)	¢1,000,000	¢C 000 000
6	\$32	\$63	\$1,000,000	\$6,000,000
N	2^{n-1}	This cell would	\$1,000,000	\$1,000,000N
1 4	2		$\psi_1,000,000$	$\varphi_1,000,0001$
		remain empty		

1. Predict when the salaries will be equal. Explain your reasoning. *In about 20 years; Explanations will vary*

2. Does Job A represent linear or exponential growth? Explain your reasoning. *Exponential; the salary is equal to some power of 2 with the power equal to the number of years.*

3. Does Job B represent linear or exponential growth? Explain your reasoning. *Linear; the rate of change is constant*

4. Write an equation to represent the salary for each job option. Job A: $s = 2^{N-1}$ Job B: s = 1,000,000N

Linear or Non-Linear?

In this activity, you will construct a scatter plot using the data set you were given and determine if the data represents a linear or non-linear relationship.

1. Identify the independent and dependent variables and write them on the top left corner of the poster.

2. Construct the scatter plot on your poster board.

3. Describe the relationship between the two variables on the top right corner of the poster, including a decision about whether your data is linear or non-linear.

4. Present your findings to the class. Include in your presentation three informative points about your data and a discussion about why the data is linear or non-linear.

Household with Television Sets (in millions)				
Year Television Sets				
1986	158			
1987	163			
1988	168			
1989	176			
1990	193			
1991	193			
1992	192			
1993	201			

Median House Price				
Year	Price			
1990	85000			
1991	88000			
1992	92000			
1993	100000			
1994	106000			
1995	115500			
1996	125000			
1997	135000			
1998	151000			
1999	160000			

Average Temperature			
Month	Тетр		
1. Jan	50		
2 Feb	54		
3 Mar	60		
4 Apr	67		
5 May	74		
6 June	80		
7 July	82		
8 Aug	81		
9 Sept	78		
10 Oct	68		
11 Nov	59		

Old Faithful geyser eruption				
Length of eruption (minutes)	Minutes between eruptions			
2	57			
2.5	62			
3	68			
3.5	75			
4	83			
4.5	89			
5	92			

Length and Weight of Whales				
Length	Weight			
(feet)	(long tons)			
40	25			
42	29			
45	34			
46	35			
50	43			
52	45			
55	51			

Wind Chill			
Wind Speed	Wind Chill		
(mph)	Fahrenheit		
0	35		
5	32		
10	22		
15	16		
20	11		
25	8		
30	6		
35	4		

World Oil Production			
Year	Barrels		
	(millions)		
1900	149		
1910	328		
1920	689		
1930	1412		
1940	2150		
1950	3803		
1960	7674		
1970	16690		
1980	21722		

Presid	Presidential Physical Fitness Awards		
	Mile – Run		
Age	Age Time (seconds)		
9	511		
10	477		
11	452		
12	431		
13	410		
14	386		

Unit 1, Activity 8, Linear or Non-Linear Rubric

RUBRIC Graphing: Linear or Non-linear?

Student Name:

CATEGORY	4	3	2	1
GRAPH				
Labeling of X axis	The X axis has a clear, neat label that describes the units used for the independent variable (e.g, days, months, participants' names).	The X axis has a clear label that describes the units used for the independent variable.	The X axis has a label.	The X axis is not labeled.
Labeling of Y axis	The Y axis has a clear, neat label that describes the units and the dependent variable (e.g, % of dog food eaten; degree of satisfaction).	The Y axis has a clear label that describes the units and the dependent variable (e.g, % of dog food eaten; degree of satisfaction).	The Y axis has a label.	The Y axis is not labeled.
Title	Title is creative and clearly relates to the problem being graphed (includes dependent and independent variable). It is printed at the top of the graph.	Title clearly relates to the problem being graphed (includes dependent and independent variable) and is printed at the top of the graph.	A title is present at the top of the graph.	A title is not present.
Accuracy of Plot	All points are plotted correctly and are easy to see. A ruler is used to neatly connect the points or make the bars, if not using a computerized graphing program.	All points are plotted correctly and are easy to see.	All points are plotted correctly.	Points are not plotted correctly OR extra points were included.
Linear or Non-linear?	Graph fits the data well and makes it easy to interpret.	Graph is adequate and does not distort the data, but interpretation of the data is somewhat difficult.	Graph distorts the data somewhat and interpretation of the data is somewhat difficult.	Graph seriously distorts the data making interpretation almost impossible.

Unit 1, Activity 8, Linear or Non-Linear Rubric

PRESENTATION	4	3	2	1
Clarity of presentation	Explanations of the project are presented in clear and detailed manner	Explanations of the project are presented in a clear manner	Explanation is a little difficult to understand, but includes critical components	Explanation is difficult to understand and includes little critical components
Accuracy of Information	All mathematical facts are reported accurately	Almost all mathematical facts are reported accurately	One fact is reported accurately	No facts were reported accurately or no facts were reported
POSTER	Exceptionally well designed, neat, and attractive. Colors that go well together are used to make the graph more readable. A ruler and graph paper (or graphing computer program) are used.	Neat and relatively attractive. A ruler and graph paper (or graphing computer program) are used to make the graph more readable.	Lines are neatly drawn, but the graph appears quite plain.	Appears messy and "thrown together" in a hurry. Lines are visibly crooked.

Creating a scatter plot with the TI-83 graphing calculator:

1) Enter data into lists:

STAT EDIT Clear lists by highlighting L1 then pressing Clear, Enter Highlight L2 press clear, enter Enter independent variable (x) in L1 Enter dependent variable (y) in L2

2) Turn on scatter plot:

 2^{nd} y=

Enter 1: On Type: scatter plot Xlist: L_1 Ylist: L_2 Mark: (any)

3) Set window to fit your data:

WINDOW Xmin and Xmax should be set to fit the data in L1 Ymin and Ymax should be set to fit the data in L2 All other entries can be set at whatever you wish

4) Look at Graph GRAPH

Unit 1, Activity 9, Understanding Data

Understanding Data

The table below gives the box score for game three of the 2003 NBA Championship series.

SAN ANTONIO SPURS						REI	BOUNE	DS			
PLAYER	POS	MIN	FGM-A	3GM-A	FTM-A	OFF	DEF	TOT	AST	PF	PTS
TONY PARKER	G	43	9-21	4-6	4-8	1	2	3	6	0	26
STEPHEN JACKSON	G	36	2-7	1-2	2-4	0	6	6	2	3	7
TIM DUNCAN	F	45	6-13	0-0	9-12	3	13	16	7	3	21
BRUCE BOWEN	F	32	0-5	0-2	0-0	1	3	4	0	3	0
DAVID ROBINSON	С	26	1-5	0-0	6-8	1	2	3	0	2	8
Emanuel Ginobili		28	3-6	0-0	2-3	2	0	2	4	2	8
Malik Rose		22	4-7	0-0	0-0	0	2	2	0	2	8
Speedy Claxton		5	2-2	0-0	0-0	0	1	1	0	1	4
Kevin Willis		3	1-1	0-0	0-0	1	0	1	0	1	2
Steve Kerr											
Danny Ferry											
Steve Smith											
TOTAL		240	28-67	5-10	23-35	9	29	38	19	17	84
			41.8%	50.0%	65.7%	Т	eam Re	ebs: 15			

NEW JERSEY NETS						REI	BOUNI	DS			
PLAYER	POS	MIN	FGM-A	3GM-A	FTM-A	OFF	DEF	TOT	AST	PF	PTS
JASON KIDD	G	42	6-19	0-5	0-0	2	1	3	11	3	12
KERRY KITTLES	G	43	8-16	3-5	2-3	1	3	4	1	2	21
KENYON MARTIN	F	42	8-18	0-1	7-8	2	9	11	0	5	23
RICHARD JEFFERSON	F	36	3-11	0-0	0-0	2	7	9	0	2	6
JASON COLLINS	С	25	0-3	0-0	0-0	4	1	5	1	6	0
Lucious Harris		22	1-6	1-2	4-4	1	0	1	3	2	7
Dikembe Mutombo		18	1-1	0-0	0-0	1	2	3	0	3	2
Rodney Rogers		11	0-3	0-0	2-2	0	2	2	0	2	2
Anthony Johnson		6	2-2	0-0	0-0	0	1	1	0	0	4
Aaron Williams		4	1-2	0-0	0-0	1	1	2	1	1	2
Tamar Slay											
Brian Scalabrine											
TOTAL		240	30-81	4-13	15-17	14	27	41	17	26	79
			37.0%	30.8%	88.2%	Т	eam Re	ebs: 10			

Source: www.	nba.com		
Key for Tab	le		
Pos	Position	3GM-A	3 point goals made – 3 point goals attempted
Min.	Minutes Played	AST	Assists
FGM-A	Field goals made-field goals attempted	PF	Personal Fouls
FTM-A	Free throws made-free throws attempted	PTS	Total Points Scored
R	Rebounds		

Use the table above to answer the following questions.

1. Which player played the most minutes of the game?

2. Who had the most assists?

3. Which team made a larger percentage of free throws?

Unit 1, Activity 9, Understanding Data

	FGM/FGA	FTM/FTA
JASON KIDD		
KERRY KITTLES		
KENYON MARTIN		
RICHARD JEFFERSON		
JASON COLLINS		
Lucious Harris		
Dikembe Mutombo		
Rodney Rogers		
Anthony Johnson		
Aaron Williams		
TONY PARKER		
STEPHEN JACKSON		
TIM DUNCAN		
BRUCE BOWEN		
DAVID ROBINSON		
Emanuel Ginobili		
Malik Rose		
Speedy Claxton		
Kevin Willis		


4. Calculate the percentage of free throws made/free throws attempted and field goals made/field goals attempted for each player. (Use the chart above to write your answers)

5. Which player(s) have/has the highest percentage? ______ Why do you think this is so?

6. Do you think that the players who attempt the most field goals are generally the players who make the most field goals?

7. Make a scatter plot showing field goals made and field goals attempted.

Identify the independent and dependent variables. Use different colors for the Spurs and Nets.



Unit 1, Activity 9, Understanding Data

8. Does the scatter plot show a negative or positive correlation?


9. Who were the four perfect shooters in the game? Circle the points on the scatter plot that represent these perfect shooters.

10. The data on the graph seems to cluster in two sections. Can you explain where they cluster and why?

11. Give three interesting facts that you notice about the scatter plot.

12. Do you think that players who get a lot of rebounds also make a lot of assists (i.e., does the number of rebounds depend on the number of assists?)

13. Construct a scatter plot of rebounds and assists.



14. Is there a relationship between the two?

Unit 1, Activity9, Understanding Data with Answers

Understanding Data

The table below gives the box score for game three of the 2003 NBA Championship series.

SAN ANTONIO SPURS						REI	BOUNE	DS			
PLAYER	POS	MIN	FGM-A	3GM-A	FTM-A	OFF	DEF	TOT	AST	PF	PTS
TONY PARKER	G	43	9-21	4-6	4-8	1	2	3	6	0	26
STEPHEN JACKSON	G	36	2-7	1-2	2-4	0	6	6	2	3	7
TIM DUNCAN	F	45	6-13	0-0	9-12	3	13	16	7	3	21
BRUCE BOWEN	F	32	0-5	0-2	0-0	1	3	4	0	3	0
DAVID ROBINSON	С	26	1-5	0-0	6-8	1	2	3	0	2	8
Emanuel Ginobili		28	3-6	0-0	2-3	2	0	2	4	2	8
Malik Rose		22	4-7	0-0	0-0	0	2	2	0	2	8
Speedy Claxton		5	2-2	0-0	0-0	0	1	1	0	1	4
Kevin Willis		3	1-1	0-0	0-0	1	0	1	0	1	2
Steve Kerr											
Danny Ferry											
Steve Smith											
TOTAL		240	28-67	5-10	23-35	9	29	38	19	17	84
			41.8%	50.0%	65.7%	T	eam Re	ebs: 15			

NEW JERSEY NETS						REI	BOUNE	DS			
PLAYER	POS	MIN	FGM-A	3GM-A	FTM-A	OFF	DEF	TOT	AST	PF	PTS
JASON KIDD	G	42	6-19	0-5	0-0	2	1	3	11	3	12
KERRY KITTLES	G	43	8-16	3-5	2-3	1	3	4	1	2	21
KENYON MARTIN	F	42	8-18	0-1	7-8	2	9	11	0	5	23
RICHARD JEFFERSON	F	36	3-11	0-0	0-0	2	7	9	0	2	6
JASON COLLINS	С	25	0-3	0-0	0-0	4	1	5	1	6	0
Lucious Harris		22	1-6	1-2	4-4	1	0	1	3	2	7
Dikembe Mutombo		18	1-1	0-0	0-0	1	2	3	0	3	2
Rodney Rogers		11	0-3	0-0	2-2	0	2	2	0	2	2
Anthony Johnson		6	2-2	0-0	0-0	0	1	1	0	0	4
Aaron Williams		4	1-2	0-0	0-0	1	1	2	1	1	2
Tamar Slay											
Brian Scalabrine											
TOTAL		240	30-81	4-13	15-17	14	27	41	17	26	79
			37.0%	30.8%	88.2%	T	eam Re	ebs: 10			

Source: www.nba.com							
Key for Table							
Pos	Position	3GM-A	3 point goals made – 3 point goals attempted				
Min.	Minutes Played	AST	Assists				
FGM-A	Field goals made-field goals attempted	PF	Personal Fouls				
FTM-A	Free throws made-free throws attempted	PTS	Total Points Scored				
R	Rebounds						

Use the table above to answer the following questions.

1. Which player played the most minutes of the game? <u>*Tim Duncan</u></u></u>*

3. Which team made a larger percentage of free throws? <u>Nets</u>

Unit 1, Activity9, Understanding Data with Answers

	FGM/FGA	FTM/FTA
JASON KIDD	32%	
KERRY KITTLES	50%	67%
KENYON MARTIN	44%	88%
RICHARD JEFFERSON	27%	
JASON COLLINS	0%	
Lucious Harris	17%	100%
Dikembe Mutombo	100%	
Rodney Rogers	0%	100%
Anthony Johnson	100%	
Aaron Williams	50%	
TONY PARKER	43%	50%
STEPHEN JACKSON	29%	50%
TIM DUNCAN	46%	75%
BRUCE BOWEN	0%	
DAVID ROBINSON	20%	75%
Emanuel Ginobili	50%	67%
Malik Rose	57%	
Speedy Claxton	100%	
Kevin Willis	100%	


4. Calculate the percentage of free throws made/free throws attempted and field goals made/field goals attempted for each player. (Use the chart above to write your answers)
5. Which player(s) have/has the highest percentage of field goals made? <u>Mutombo, A.</u> Johnson, Claxton, Willis_

Why do you think this is so? *They did not have as many attempts*

6. Do you think that the players who attempt the most field goals are generally the players who make the most field goals? <u>You can't make goals if you don't attempt them</u>.

7. Make a scatter plot showing field goals made and field goals attempted.

Identify the independent (*attempted*) and dependent variables.(*made*)Use different colors for the Spurs and Nets.



Unit 1, Activity9, Understanding Data with Answers

8. Does the scatter plot show a negative or positive correlation? *positive*

9. Who were the four perfect shooters in the game? Circle the points on the scatter plot that represent these perfect shooters. *Mutombo, A. Johnson, Claxton, Willis*_____


10. The data on the graph seems to cluster in two sections. Can you explain where they cluster and why?

The cluster at the highest point represents the players who played more minutes. The lower cluster is the group that played fewer minutes.

11. Give three interesting facts that you notice about the scatter plot. *Answers will vary*

12. Do you think that players who get a lot of rebounds also make a lot of assists (i.e. does the number of rebounds depend on the number of assists)? *Students' opinions will vary*

13. Construct a scatter plot of rebounds and assists.



14. Is there a relationship between the two? no