

## HEDIS 2020 CODING GUIDE

In addition to the coding guide, care gap reports can be produced to identify members with gaps in care according to HEDIS specifications. The Health Plan provider engagement representatives can distribute gap reports to primary care physicians. Clinical documentation to support HEDIS measures can be submitted via fax to 1.304.433.8208 or by contacting your provider engagement representative.

| Code   | Measure Title  | Product Line             | Description   | Coding<br>(Identifying Numerator)  | Category II<br>Codes-CPT |
|--------|--|--------------------------|---|--|--------------------------|
| PREVEN | NTION AND SCREENIN   | IG                       |   |  |                          |
| АВА    | Adult BMI<br>Assessment  | ALL                      | Percentage of members 18-74 who had<br>their BMI and weight documented at an<br>outpatient visit within the past 2 years<br>*For members younger than 20 years of<br>age documentation must be a BMI<br>percentile with height and weight                                   | <ul> <li>BMI: ICD-10: Z68.1, Z68.20-68.39, Z68.41-68.45</li> <li>BMI Percentile: ICD-10: Z68.51-68.54</li> </ul>   |                          |
| wcc    | Weight Assessment<br>and Counseling for<br>Nutrition and<br>Physical Activity for<br>Children and<br>Adolescents | Commercial<br>& Medicaid | Percentage of members 3-17 who had an<br>outpatient visit with a PCP or OB/GYN<br>which included evidence of BMI<br>documentation with corresponding height<br>& weight, counseling for nutrition and/or<br>counseling for physical activity during the<br>measurement year | BMI Percentile: ICD-10: Z68.51-         68.54         Nutrition Counseling: CPT: 97802-         97804; HCPCS: S9470, S9452, S9449,         G0270, G0271, G0447; ICD-10:         Z71.3         Physical Activity Counseling:         HCPCS: S9451, G0447; ICD-10:         Z02.5, Z71.82 |                          |

| Code  | Measure Title                    | Product Line             | Description   | Coding<br>(Identifying Numerator)   | Category II<br>Codes-CPT |
|-------|----------------------------------|--------------------------|---|---|--------------------------|
| PREVE | NTION AND SCREENIN               | IG, CONT.                |   |   |                          |
| CIS   | Childhood<br>Immunization Status | Commercial<br>& Medicaid | Percentage of children two years of age<br>with appropriate childhood immunizations<br>including at least:<br>• 4 DTAP<br>• 3 IPV<br>• 1 MMR<br>• 3 HIB<br>• 3 Hep B<br>• 1 VZV<br>• 4 PCV<br>• 1 Hep A<br>• 2-3 dose Rotavirus vaccines<br>• At least 2 Influenza vaccines   | DTaP: CPT: 90698, 90700, 90721,<br>90723<br>IPV: CPT: 90698, 90713, 90723<br>MMR: CPT: 90707, 90710<br>Hib: CPT: 90644-90648, 90698,<br>90721, 90748<br>Hepatitis B: CPT: 90723, 90740,<br>90744, 90747, 90748; HCPCS:<br>G0010<br>VZV: CPT: 90710, 90716<br>Pneumococcal Conjugate: CPT:<br>90670 HCPCS: G0009<br>Hepatitis A: CPT: 90633<br>Rotavirus (2 dose schedule): CPT:<br>90681<br>Rotavirus (3 dose schedule): CPT:<br>90681<br>Rotavirus (3 dose schedule): CPT:<br>90680<br>Influenza: CPT: 90655, 90657, 90661,<br>90662, 90673, 90685-90688; HCPCS:<br>G0008<br>Exclusion (Anaphylactic<br>Reaction): ICD-10: T80.52XA,<br>T80.52XD, T80.52XS |                          |
| IMA   | Immunizations for<br>Adolescents | Commercial<br>& Medicaid | Percentage of adolescents 13 years of<br>age with appropriate immunizations<br>including at least:<br>• 1 meningococcal vaccine between 11th<br>and 13th birthdays<br>• 1 Tdap between 10th and 13th birthdays;<br>• 2 HPV vaccines at least 146 days apart <u>or</u><br>3 HPV vaccines on different dates of<br>service between 9th and 13th birthdays | Meningococcal: CPT: 90734<br>Tdap: CPT: 90715<br>HPV: CPT: 90649, 90650, 90651<br>Exclusion (Anaphylactic<br>Reaction): ICD-10: T80.52XA,<br>T80.52XD, T80.52XS   |                          |

| Code  | Measure Title                   | Product Line             | Description  | Coding<br>(Identifying Numerator)  | Category II<br>Codes-CPT |  |  |  |
|-------|---------------------------------|--------------------------|--|--|--------------------------|--|--|--|
| PREVE | PREVENTION AND SCREENING, CONT. |                          |  |  |                          |  |  |  |
| BCS   | Breast Cancer<br>Screening      | ALL                      | Percentage of women 50-74 years of age<br>who had a mammogram any time on or<br>between Oct 1 two years prior to the<br>measurement year and Dec 31 of the<br>measurement year   | Mammography: CPT: 77055-77057,<br>77061-77063, 77065-77067; HCPCS:<br>G0202, G0204, G0206<br>Exclusion (Bilateral Mastectomy):<br>ICD-10: OHTV0ZZ, ICD-9: 85.42,<br>85.44, 85.46, 85.48  |                          |  |  |  |
| CCS   | Cervical Cancer<br>Screening    | Commercial<br>& Medicaid | Percentage of women 21-64 years of age<br>who had a Pap test within the past 3 years<br>or women 30-64 who had cervical<br>cytology/high-risk human papillomavirus<br>(hrHPV) cotesting within the past 5 years  | Cervical Cytology: CPT: 88141-<br>88143, 88147, 88148, 88150, 88152-<br>88154, 88164-88167, 88174, 88175<br>HPV: CPT: 87620-87622, 87624-<br>87625 HCPCS: G0123, G0124,<br>G0141, G0143-G0145, G0147,<br>G0148, P3000, P3001, Q0091<br>Exclusion (Hysterectomy): CPT:<br>51925, 56308, 57540, 57545, 57550,<br>57555, 57556, 58150, 58152, 58200,<br>58210, 58240, 5860, 58262, 58263,<br>58267, 58270, 58275, 58280, 58285,<br>58290-58294, 58548, 58550, 58552-<br>58554, 58570-58575, 58951,58953,<br>58954,58956, 59135  |                          |  |  |  |
| COL   | Colorectal Cancer<br>Screening  | Commercial<br>& Medicare | Percentage of members 50-75 years of<br>age who had appropriate screening for<br>colorectal cancer including at least <u>one</u><br>of the following:<br>• FOBT during the measurement year<br>• Flexible sigmoidoscopy within the past 5<br>years<br>• Colonoscopy within the past 10 years<br>• CT colonography within the past 5 years<br>• FIT-DNA within the past 3 years | FOBT: CPT: 82270, 82274; HCPCS:         G0328         FIT-DNA: CPT: 81528; HCPCS:         G0464         CT Colonography: CPT: 74261-         74263         Flexible Sigmoidoscopy: CPT:         45330-45335, 45337-45342, 45345-         45347, 45349, 45350; HCPCS:         G0104         Colonoscopy: CPT: 44388-44394,         44397, 44401-44408, 45355, 45378-         45393, 45398; HCPCS: G0105,         G0121         Exclusion (Colorectal Cancer):         HCPCS: G0213, G0214, G0215,         G0231; ICD-10: C18.0-C18.9, C19,         C20, C21.8, C78.5, Z85.038, Z85.048         (Total Colectomy): CPT: 44150-         44153, 44155-44158, 44210-44212 |                          |  |  |  |

| Code   | Measure Title   | Product Line   | Description   | Coding<br>(Identifying Numerator)   | Category II<br>Codes-CPT      |
|--------|---|--|---|---|-------------------------------|
| PREVE  | NTION AND SCREENIN  | IG, CONT.  |   |   |                               |
| CHL    | Chlamydia<br>Screening in<br>Women  | Commercial<br>& Medicaid   | Sexually active women 16-24 with annual chlamydia screening   | <b>CPT:</b> 87110, 87270, 87320, 87490-<br>87492, 87810   |                               |
|        |   |  | Percentage of adults 66 years and older who had <u>each</u> of the following during   | <b>Advance Care Planning: CPT:</b><br>99483, 99497<br><b>HCPCS:</b> \$0257  | 1123F, 1124F,<br>1157F, 1158F |
| СОА    | Care for Older<br>Adults  | Medicare<br>SNP  | <ul> <li>the measurement year:</li> <li>Advance care planning</li> <li>Medication review (includes a</li> </ul>   | Medication Review: CPT: 90863,<br>99483, 99605, 99606<br>Medication List: HCPCS: G8427  | 1160F<br>1159F                |
|        | Medication review (include<br>medication list and a review     Functional status assessment | <ul> <li>Medication list and a review)</li> <li>Functional status assessment</li> <li>Pain assessment</li> </ul> | Functional Status Assessment:<br>CPT: 99483; HCPCS: G0438,<br>G0439   | 1170F   |                               |
|        |   |  |   | Pain Assessment:  | 1125F, 1126F                  |
| RESPIR | ATORY CONDITIONS  |  |   |   |                               |
| CWP    | Appropriate<br>Testing for<br>Pharyngitis   | Commercial,<br>Medicaid, &<br>Medicare   | Percentage of members 3 years or older<br>diagnosed with pharyngitis, prescribed<br>an antibiotic, and tested for strep   | <b>CPT:</b> 87070, 87071, 87081, 87430, 87650-87652, 87880  |                               |
| PCE    | Pharmacotherapy<br>Management of<br>COPD<br>Exacerbation                                    | ALL  | Members dispensed systemic<br>corticosteroid within 14 days or<br>bronchodilator within 30 days of an<br>acute inpatient discharge or ED<br>encounter with a principal diagnosis of<br>COPD | Dispensed Systemic<br>Corticosteroid and<br>Bronchodilator  |                               |
| мма    | Medication<br>Management for<br>People with<br>Asthma                                       | Commercial<br>& Medicaid   | Percentage of members 5-64 years who<br>remain on an asthma controller for at<br>least 75% of their treatment period  | Frequency of Prescribed<br>Preferred Asthma Medications:<br>antiasthmatic combinations,<br>antibody inhibitor, inhaled<br>steroid combinations, inhaled<br>corticosteroids, leukotriene<br>modifiers, mast cell stabilizers,<br>methylxanthines |                               |

| Code    | Measure Title   | Product Line             | Description   | Coding<br>(Identifying Numerator)  | Category II<br>Codes-CPT  |
|---------|---|--------------------------|---|--|---|
| RESPIRA | ATORY CONDITIONS,   | CONT.                    |   |  |   |
| AMR     | Asthma<br>Medication Ratio  | Commercial<br>& Medicaid | Percentage of members 5-64 years of<br>age with asthma who had a ratio of<br>controller medications to total asthma<br>medications of .5 or greater   | Asthma Controller and Reliever<br>Medications: Antiasthmatic<br>combinations, antibody<br>inhibitors, inhaled steroid<br>combinations, inhaled<br>corticosteroids, leukotriene<br>modifiers, mast cell stabilizers,<br>methylxanthines; short acting,<br>inhaled beta-2 agonists |   |
| CARDI   | OVASCULAR   |                          |   |  |   |
| Свр     | Controlling High<br>Blood Pressure                                  | ALL                      | Percentage of members 18-85 years of<br>age with a diagnosis of hypertension<br>and whose blood pressure was<br>controlled (<140/90 mm Hg) during the<br>measurement year   |  | <b>Systolic:</b> 3074F,<br>3075F, 3077F<br><b>Diastolic:</b> 3078F,<br>3079F, 3080F |
| РВН     | Persistence of<br>Beta-Blocker<br>Treatment After a<br>Heart Attack | ALL                      | Percentage of members 18 years or<br>older who were discharged with a<br>diagnosis of acute myocardial infarction<br>and received a beta-blocker treatment<br>for 6 months after discharge  | <b>Beta Blocker Medications:</b><br>Noncardioselective beta-<br>blockers, Cardioselective beta-<br>blockers, Antihypertensive<br>combinations  |   |
| SPC     | Statin Therapy for<br>Patients with<br>Cardiovascular<br>Disease    | ALL                      | The percentage of males 21-75 years of<br>age and females 40-75 years of age<br>with clinical atherosclerotic<br>cardiovascular disease (ASCVD) and<br>were dispensed at least one high or<br>moderate intensity statin medication<br>during the measurement year and<br>remained on medication for at least<br>80% of the treatment period | High intensity statin therapy<br>medications<br>Moderate intensity statin therapy<br>medications   |   |

| Code   | Measure Title                                   | Product Line | Description   | Coding<br>(Identifying Numerator)   | Category II<br>Codes-CPT  |
|--------|---|--------------|---|---|---|
| DIABET | ES  |              |   |   |   |
|        |   |              |   | HbA1c Tests: CPT: 83036, 83037  | 3044F, 3045F,<br>3046F  |
|        |   |              |   | Nephropathy Screening: CPT:<br>81000-81003, 81005, 82042-82044,<br>84156  | 3060F, 3061F,<br>3062F  |
| CDC    | Comprehensive<br>Diabetes Care                  | ALL          | The percentage of members 18-75 years<br>of age with diabetes who had <u>each</u> of<br>the following during the measurement<br>year:<br>• HbA1c (control <8.0%)<br>• Nephropathy screening<br>• Retinal eye exam   | <b>Eye Exams: CPT:</b> 67028, 67030,<br>67031, 67036, 67039-67043, 67101,<br>67105, 67107, 67108, 67110,<br>67113, 67121, 67141, 67145,<br>67208, 67210, 67218, 67220,<br>67221, 67227, 67228, 92002,<br>92004, 92012, 92014, 92018,<br>92019, 92134, 92225-92228, 92230,<br>92235, 92240, 92250, 92260,<br>99203-99205, 99213-99215, 99242-<br>99245 HCPCS: S0620, S0621,<br>S3000 | 2022F, 2024F,<br>2026F, 3072F   |
|        |   |              | Blood pressure control (<140/90)  | Blood pressure control:Systolic:<br>3075F, 30<br>Diastolic:   | Systolic: 3074F,<br>3075F, 3077F<br>Diastolic: 3078F,<br>3079F, 3080F |
|        |   |              |   |   |   |
| SPD    | Statin Therapy for<br>Patients with<br>Diabetes | ALL          | The percentage of diabetic members<br>40-75 years of age who do <b>not</b> have<br>clinical atherosclerotic cardiovascular<br>disease (ASCVD) and were dispensed at<br>least one statin medication during the<br>measurement year and remained on<br>medication for at least 80% of the<br>treatment period | High intensity statin therapy<br>medications<br>Moderate intensity statin therapy<br>medications<br>Low intensity statin therapy<br>medications   |   |

| Code  | Measure Title  | Product Line | Description   | Coding<br>(Identifying Numerator)  | Category II<br>Codes-CPT |
|-------|--|--------------|---|--|--------------------------|
| MUSCI | JLOSKELETAL  |              |   |  |                          |
| ART   | Disease-Modifying<br>Anti-Rheumatic<br>Drug Therapy for<br>Rheumatoid<br>Arthritis | Medicare     | Percentage of members 18 years of age<br>and older who were diagnosed with<br>rheumatoid arthritis and who were<br>dispensed at least one ambulatory<br>prescription for a disease-modifying<br>anti-rheumatic drug (DMARD) during<br>the measurement year  | A dispensed DMARD medication<br><b>or HCPCS:</b> J0129, J0135, J0717,<br>J1438, J1600, J1602, J1745, J3262,<br>J7502, J7515-J7518, J9250, J9260,<br>J9310, Q5102-Q5104   |                          |
| омw   | Osteoporosis<br>Management in<br>Women Who Had<br>Fracture                         | Medicare     | Percentage of women 67-85 years of<br>age who suffered a fracture and<br>received appropriate osteoporosis<br>treatment as indicated by <u>one</u> of the<br>following:<br>• Bone Mineral Density Test within 6<br>months of the fracture or during the<br>inpatient stay if fracture occurred during<br>hospitalization<br>• Osteoporosis therapy within 6 months<br>of the fracture<br>• Long-acting osteoporosis therapy<br>during the inpatient stay if fracture<br>occurred during hospitalization<br>• A dispensed prescription to treat<br>osteoporosis within 6 months of the<br>fracture | <b>CPT:</b> 76977, 77078, 77080-77082,<br>77085, 77086<br><b>HCPCS:</b> G0130, J0630, J0897,<br>J1740, J3110, J3489, J2794, J2680,<br>J2426, J2358, J1631, J0401<br><b>or</b> Biphosphonates or other<br>approved agents                                       |                          |
| BEHAV | IORAL HEALTH   |              |   |  |                          |
| АММ   | Antidepressant<br>Medication<br>Management   | ALL          | Percentage of members 18 years or<br>older diagnosed with depression who<br>remained on antidepressant medication<br>for at least 6 months<br>**This measure is a 12 month window<br>starting on May 1 of the year prior to the<br>measurement year and ending on April<br>30 of the measurement year.  | Miscellaneous antidepressants,<br>Monoamine oxidase inhibitors,<br>Phenylpiperazine<br>antidepressants,<br>Psychotherapeutic<br>combinations, SNRI<br>antidepressants, SSRI<br>antidepressants, Tetracycline<br>antidepressants, Tricycline<br>antidepressants |                          |

| Code  | Measure Title  | Product Line             | Description   | Coding<br>(Identifying Numerator)  | Category II<br>Codes-CPT |
|-------|--|--------------------------|---|--|--------------------------|
| BEHAV | IORAL HEALTH, CONT   |                          |   |  |                          |
| ADD   | Follow-Up Care for<br>Children Prescribed<br>ADHD Medication | Commercial<br>& Medicaid | Percentage of children 6-12 years of age<br>with newly prescribed ADHD medication<br>who received the appropriate follow-up<br>visits and medication. Appropriate visits<br>include an initiation visit 30-day post<br>medication and two visits within 9 months<br>of the initiation visit.<br>**This measure is a 12 month window<br>starting on March 1 of the year prior to the<br>measurement year and ending the last<br>day of February of the measurement year. | Initiation Visit: CPT: 90791, 90792, 90832-<br>90834, 90836-90840, 90845, 90847,<br>90849, 90853, 90875, 90876, 99221-<br>99223, 99231-99233, 99238, 99239,<br>99251-99255, 98960-98962, 99078,<br>99201-99205, 99211-99215, 99241-99245,<br>99341-99345, 99347-99350, 99381-99387,<br>99391-99397, 99401-99404, 99411, 99412,<br>99483, 99510; HCPCS: G0155, G0176,<br>G0177, G0409, G0463, H0002, H0004,<br>H0031, H0034, H0036, H0037, H0039,<br>H0040, H2000, H2010, H2011, H2013-<br>H2020, M0064<br>Maintenance Visit: CPT: 90791, 90792,<br>90832-90834, 90836-90840, 90845, 90847,<br>90849, 90853, 90875, 90876, 96150-<br>96154, 98966-99443, 99221-99223,<br>99231-99233, 99238, 99239, 99251-99255,<br>98960-98962, 99078, 99201-99205,<br>99211-99215, 99241-99245, 99341-99345,<br>99347-99350, 99381-99387, 99391-99397,<br>99401-99404, 99411, 99412, 99483,<br>99510; HCPCS: G0155, G0176, G0177,<br>G0409, G0463, H0002, H0004, H0031,<br>H0034, H0036, H0037, H0039, H0040,<br>H2000, H2010, H2011, H2013-H2020,<br>M0064 |                          |
| FUH   | Follow-Up After<br>Hospitalization for<br>Mental Illness     | ALL                      | Percentage of discharges for members 6<br>years or older who were hospitalized for<br>treatment of a mental health disorder and<br>received appropriate follow-up visits within<br>7 and 30 days during the measurement<br>year   | <b>CPT:</b> 90791, 90792, 90832-90834, 90836-<br>90840, 90845, 90847, 90849, 90853,<br>90875, 90876, 96150-96154, 99221-<br>99223, 99231-99233, 99238, 99239,<br>99251-99255, 98960-98962, 99078,<br>99201-99205, 99211-99215, 99241-<br>99245, 99341-99345, 99347-99350,<br>99381-99387, 99391-99397, 99401-<br>99404, 99411, 99412, 99483, 99510;<br><b>HCPCS:</b> G0155, G0176, G0177, G0409,<br>G0463, H0002, H0004, H0031, H0034,<br>H0036, H0037, H0039, H0040, H2000,<br>H2010, H2011, H2013-H2020, M0064   |                          |

| Code  | Measure Title   | Product Line             | Description   | Coding<br>(Identifying Numerator)   | Category II<br>Codes-CPT |
|-------|---|--------------------------|---|---|--------------------------|
| BEHAV | IORAL HEALTH, CONT  | •                        |   |   |                          |
| FUI   | Follow-up After High<br>Intensity Care for<br>Substance Use<br>Disorder         | ALL                      | Percentage of members 13 years of age<br>and older who receive follow-up<br>treatment within 7 and 30 days following<br>acute inpatient hospitalization, residential<br>treatment, or detoxification visit for<br>substance use disorder during the<br>measurement year           | <b>CPT</b> 90791, 90792, 90832-90834,<br>90836-90840, 90845, 90847, 90849,<br>90853, 90875, 90876, 98960-98962,<br>98966-98969, 99078, 99201-99205,<br>99211-99215, 99221-99223, 99231-<br>99233, 99238, 99239, 99241-99245,<br>99251-99255, 99341-99345, 99347-<br>99350, 99384-99387, 99394-99397,<br>99401-99404, 99408, 99409, 99411,<br>99412, 99441-99444, 99483, 99510;<br><b>HCPCS:</b> H0020, H0033, J0570-J0575,<br>J2315, Q9991, Q9992, S0109,<br>G0155, G0176, G0177, G0396,<br>G0397, G0409-G0411, G0443,<br>G0463, H0001, H0002, H0004,<br>H0005, H0007, H0015-H0019, H0022,<br>H0031, H0034-H0037, H0039, H0040,<br>H0047, H2000, H2001, H2010-H2020,<br>H2035, H2036, M0064, S0201, S9480,<br>S9484, S9485, T1006, T1012, T1015,<br>T2048 |                          |
| АРМ   | Metabolic<br>Monitoring for<br>Children and<br>Adolescents on<br>Antipsychotics | Commercial<br>& Medicaid | Percentage of children and adolescents<br>1-17 years of age who had two or more<br>antipsychotic prescriptions and had<br>metabolic testing including at least <u>one</u> of<br>the following during the measurement<br>year:<br>• Blood glucose testing<br>• Cholesterol testing | <b>CPT:</b> 80047, 80048, 80050, 80053, 80069, 82947, 82950, 82951, 83036, 83037, 82465, 83718, 84478, 80061, 83700, 83701, 83704, 83721  |                          |
| POD   | Pharmacotherapy<br>for Opioid Use<br>Disorder                                   | ALL                      | Percentage of new opioid use disorder<br>pharmacotherapy events with OUD<br>pharmacotherapy for 180 days or more<br>days among members age 16 and older<br>with this diagnosis  | Naltrexone oral medications,<br>Naltrexone Injection medications<br>list, Buprenorphine Oral<br>medications list, Buprenorphine<br>Implant Medications list,<br>Buprenorphine Naloxone<br>Medication list   |                          |

| Code  | Measure Title  | Product Line | Description  | Coding<br>(Identifying Numerator)  | Category II<br>Codes-CPT |
|-------|--|--------------|--|--|--------------------------|
| BEHAV | IORAL HEALTH, CONT   |              |  |  |                          |
| FUM   | Follow up After<br>Emergency<br>Department Visit for<br>Mental Illness   | ALL          | Percentage of ED visits for members 6<br>years of age and older with a principal<br>diagnosis of mental illness or intentional<br>self-harm who had a follow-up visit for<br>mental illness  | Follow-up visit for mental illness<br>(follow-up within 7 days of<br>discharge; follow-up within 30 days<br>of discharge)  |                          |
| FUA   | Follow up After<br>Emergency<br>Department Visit for<br>Alcohol and Other<br>Drug Abuse or<br>Dependence           | ALL          | Percentage of emergency department<br>visits for members 13 years of age or older<br>with a principal diagnosis of alcohol or<br>other drug dependence who had a follow<br>up for AOD during the measurement year                      | Follow-up visit for AOD (follow-up<br>within 7 days of discharge; follow-<br>up within 30 days of discharge)   |                          |
| SSD   | Diabetes Screening<br>for People with<br>Schizophrenia or<br>Bipolar Who Are<br>Using Antipsychotic<br>Medications | Medicaid     | Percentage of members 18-64 years of<br>age with schizophrenia, schizoaffective<br>disorder or bipolar disorder, who were<br>dispensed an antipsychotic medication<br>and had a diabetes screening test during<br>the measurement year | HbA1c Tests: CPT: 83036, 83037<br>Antipsychotic Medication:<br>Aripiprazole,<br>Asenapine,Brexpiprazole,<br>Cariprazine, Clozapine,<br>Haloperidol, lioperidone, Loxapine,<br>Lurisadone, Molindone,<br>Olanzapine, Paliperidone,<br>Quetiapine, Quetiapine fumerate,<br>Risperidone, Ziprasidone,<br>Chlorpromazine, Fluphenazine,<br>Perphenazine, Prochlorperazine,<br>Thioridazine, Trifluoperazine,<br>Amitriptyline-perphenazine,<br>Thiothizene, Aripiprazole,<br>Fluenazine decanoate,<br>Haloperidol decanoate,<br>Olanzapine, Paliperidone<br>palmitate, Risperidone |                          |

| Code  | Measure Title   | Product Line | Description   | Coding<br>(Identifying Numerator)   | Category II<br>Codes-CPT |
|-------|---|--------------|---|---|--------------------------|
| BEHAV | IORAL HEALTH, CON   | r.           |   |   |                          |
| SAA   | Adherence to<br>Antipsychotic<br>Medications for<br>Individuals with<br>Schizophrenia | ALL          | Percentage of members 18 years and<br>older during the measurement year with<br>schizophrenia or schizoaffective disorder<br>who were dispensed and remained on<br>an antipsychotic medication for at least<br>80% of their treatment period  | Dementia Medication:<br>Donepezil, Galantamine,<br>Rivastigmine, Memantime<br>Antipsychotic Medications:<br>Aripiprazole, Asenapine,<br>Brexpiprazole, Cariprazine,<br>Clozapine, Haloperidol,<br>lioperidone, Loxapine,<br>Lurisadone, Molindone,<br>Olanzapine, Paliperidone,<br>Quetiapine, Quetiapine<br>fumarate, Risperidone,<br>Ziprasidone, Chlorpromazine,<br>Fluphenazine, Perphenazine,<br>Prochlorperazine, Thioridazine,<br>Trifluoperazine, Amitriptyline-<br>perphenazine, Thiothixene<br>Long acting injections:<br>Risperidone, Aripiprazole,<br>Fluphenazine decanoate,<br>Haloperidol deconoate,<br>Olanzapine Paliperidone<br>palmitate |                          |
| MEDIC | ATION MANAGEMEN   | T            |   |   |                          |
| TRC   | Transitions of Care   | Medicare     | Percentage of discharges during the<br>measurement year for members 18<br>years of age and older who had <u>each</u><br>of the following:<br>• Notification of inpatient admission (on<br>day of admission or following day)<br>• Receipt of discharge information (on<br>day of discharge or following day)<br>• Patient engagement after inpatient<br>discharge (within 30 days after<br>discharge)<br>• Medication reconciliation post-<br>discharge (within 30 days after | Patient Engagement: CPT: 99201-         99205, 99211-99215, 99241-99245,         99341-99345, 99347-99350, 99381-         99387, 99391-99397, 99401-99404,         99411, 99412, 99429, 99455,         99456, 99483, 98966-98968,         99441-99443; HCPCS: H0402,         G0438, G0439, G0463, T1015         Medication Reconciliation:         99483, 99495, 99496  | 1111F                    |

| Code  | Measure Title   | Product Line | Description  | Coding<br>(Identifying Numerator)   | Category II<br>Codes-CPT |
|-------|---|--------------|--|---|--------------------------|
| MEDIC | AL MANAGEMENT, C  | ONT.         |  |   |                          |
| MRP   | Medication<br>Reconciliation<br>Post-Discharge  | Medicare     | Percentage of discharges during the<br>measurement year for members 18<br>years of age and older for whom<br>medications were reconciled on or<br>within 30 days of discharge  | <b>CPT:</b> 99483, 99495, 99496   | 1111F                    |
| FMC   | Follow up After<br>Emergency<br>Department Visit<br>for People with<br>Multiple High-Risk<br>Chronic Conditions | Medicare     | Percentage of members 18 years and<br>older with multiple high-risk chronic<br>conditions who had a follow-up service<br>within 7 days of an ED visit during the<br>measurement year   | Follow-up visit within 7 days of ED<br>visit  |                          |
| OVERU | ISE/APPROPRIATENES  | S            |  |   |                          |
| PSA   | Non-<br>Recommended<br>PSA-Based<br>Screening in Older<br>Men   | Medicare     | Percentage of men 70 years and older<br>during the measurement year who were<br>screened unnecessarily for prostate<br>cancer using prostate-specific antigen<br>(PSA)-based screening<br>**A lower rate indicates better<br>performance   | CPT: 84152-84154; HCPCS: G0103  |                          |
| URI   | Appropriate<br>Treatment for<br>Upper Respiratory<br>Infection  | ALL          | Percentage of children 3 months of age<br>and older diagnosed with upper<br>respiratory infection diagnosis and <b>not</b><br>dispensed an antibiotic**This measure<br>has a 12-month window starting July 1 of<br>the year prior to the measurement year<br>and ending June 30 of the<br>measurement year | Antibiotic Prescriptions:<br>Aminopenicillins, Beta-lactamase<br>inhibitors, first generation<br>cephalosporins, folate<br>antagonist, lincomycin<br>derivatives, macrolides,<br>miscellaneous antibiotics, natural<br>penicillins, penicillinase-resistant<br>penicillins, quinolones, second<br>generation cephalosporins,<br>sulfonamides, tetracyclines, third<br>generation cephalosporins |                          |

| Code  | Measure Title   | Product Line             | Description  | Coding<br>(Identifying Numerator)   | Category II<br>Codes-CPT |  |  |
|-------|---|--------------------------|--|---|--------------------------|--|--|
| OVERU | OVERUSE/APPROPRIATENESS, CONT.                                  |                          |  |   |                          |  |  |
| AAB   | Avoidance of<br>Antibiotic<br>Treatment for<br>Acute Bronchitis | ALL                      | Percentage of members 3 months and<br>older diagnosed with acute bronchitis<br>who were <b>not</b> dispensed an antibiotic<br>**This measure has a 12-month<br>window starting July 1 of the year prior<br>to the measurement year and ending<br>June 30 of the measurement year | Antibiotic Prescriptions:<br>Aminoglycosides,<br>aminopenicillins,<br>antipseudomonal penicillins,<br>beta-lactamase inhibitors, first<br>generation cephalosporins,<br>fourth generation<br>cephalosporins, ketolides,<br>lincomycin derivatives,<br>macrolides, miscellaneous<br>antibiotics, natural penicillins,<br>penicillinase-resistant penicillins,<br>quinolones, rifamycin derivatives,<br>second generation<br>cephalosporin, sulfonamides,<br>tetracyclines, third generation<br>cephalosporins, urinary anti-<br>infectives |                          |  |  |
| LBP   | Use of Imaging<br>Studies for Low<br>Back Pain                  | Commercial<br>& Medicaid | Percentage of members with a<br>primary diagnosis of low back pain<br>during the measurement year who did<br><b>not</b> have an imaging study within 28<br>days of diagnosis   | <b>CPT:</b> 72020, 72052, 72100, 72110, 72114, 72120, 72131-72133, 72141, 72142, 72146-72149, 72156, 72158, 72200, 72202, 72220   |                          |  |  |
| DAE   | Use of High-Risk<br>Medications in<br>Older Adults              | Medicare                 | The percentage of members over 66<br>years of age who had at least two<br>dispensing events for the same high-<br>risk medication during the<br>measurement year<br>**A lower rate indicates better<br>performance   | Members who had two<br>dispensing events for the same<br>high-risk medication   |                          |  |  |

| Code                           | Measure Title   | Product Line | Description   | Coding<br>(Identifying Numerator)   | Category II<br>Codes-CPT |  |
|--------------------------------|---|--------------|---|---|--------------------------|--|
| OVERUSE/APPROPRIATENESS, CONT. |   |              |   |   |                          |  |
| DDE                            | Potentially Harmful<br>Drug-Disease<br>Interaction in Older<br>Adults | Medicare     | Percentage of members 65 years or<br>older who have evidence of an<br>underlying disease, condition or heath<br>concern who were dispensed an<br>ambulatory prescription for a<br>contraindicated medication during the<br>measurement year<br>**A lower rate indicates better<br>performance | Members identified with falls or<br>hip fractures: Dispensed an<br>ambulatory prescription for an<br>anticonvulsant, antipsychotic,<br>benzodiazepine,<br>nonbenzodiazepine hypnotic,<br>SSRI, SNRI, or tricylic<br>antidepressant<br>Members diagnosed with<br>dementia or a dispensed<br>dementia medication: Dispensed<br>an ambulatory prescription for<br>an antipsychotic,<br>benzodiazepine,<br>nonbenzodiazepine hypnotic,<br>tricylic antidepressant, or<br>anticholinergic agent<br>Members diagnosed with<br>chronic renal failure: Dispensed<br>an ambulatory prescription for a<br>nonaspirin NSAID or Cox-2 NSAID |                          |  |
| HDO                            | Use of Opioids at<br>High Dose  | ALL          | Proportion of members 18 years and<br>older who received prescription opioids<br>at a high dose for >15 days during the<br>measurement year.  | Members who received<br>prescription opioids at high<br>dosage for ≥ 15 days  |                          |  |
| UOP                            | Use of Opioids from<br>Multiple Providers                             | ALL          | Proportion of members 18 years and<br>older, receiving prescription opioids for<br>≥15 days during measurement year who<br>received opioids from multiple providers   | Receiving prescription opioids at<br>high dosage for ≥ 15 days,<br>received from mulitple providers<br>1. Multiple prescribers (four or<br>more prescribers) 2. Multiple<br>pharmacies (four or more<br>pharmacies) 3. Multiple<br>prescribers & pharmacies (both<br>four or more prescribers <b>and</b> four<br>or more pharmacies)  |                          |  |

| Code                           | Measure Title   | Product Line             | Description   | Coding<br>(Identifying Numerator)  | Category II<br>Codes-CPT |  |
|--------------------------------|---|--------------------------|---|--|--------------------------|--|
| OVERUSE/APPROPRIATENESS, CONT. |   |                          |   |  |                          |  |
| COU                            | Risk of Continued<br>Opioid Use   | ALL                      | Percentage of members 18 years or<br>older who have a new episode of<br>opioid use during the measurement<br>year that puts them at risk for continued<br>opioid use  | Have new episode of opioid use<br>that puts them at risk for<br>continued opioid use. 1.<br>Percentage of members whose<br>new episode of opioid use lasts<br>at least 15 days in a 30-day<br>period 2. Percentage of<br>members whose new episode of<br>opioid use last at least 31 days in<br>62-day period  |                          |  |
| ACCES                          | S/AVAILABILITY OF C   |                          |   |  |                          |  |
| САР                            | Children and<br>Adolescents;<br>Access to Primary<br>Care Practitioners | Commercial<br>& Medicaid | Percentage of members 1-6 years of<br>age who had a visit with a PCP during<br>the measurement year. Children and<br>adolescents 7-19 years of age who had<br>a visit with a PCP during the<br>measurement year or the year prior to<br>the measurement year. | <b>CPT:</b> 99201-99205, 99211-99215,<br>99241-99245, 99341-99345, 99347-<br>99350, 99381-99387, 99391-99397,<br>99401-99404, 99411, 99412, 99429,<br>99483 <b>HCPCS:</b> G0402, G0438,<br>G0439, G0463, T1015 <b>ICD-10</b> :<br>Z00.00, Z00.01, Z00.121, Z00129,<br>Z00.3, Z00.5, Z00.8, Z02.0Z02.6,<br>Z02.71, Z02.79, Z02.81-Z02.83,<br>Z02.89, Z02.9, 76.1, Z76.2   |                          |  |
| ААР                            | Adults' Access to<br>Preventive/<br>Ambulatory Health<br>Services       | ALL                      | The percentage of members 20 years or<br>older who had an ambulatory or<br>preventive care visit during:<br>• the measurement year (Medicaid and<br>Medicare)<br>or<br>• the measurement year or the two<br>years prior (Commercial)                          | <b>CPT:</b> 98966-98968, 99441-99443,<br>92002, 92004, 92012, 92014,<br>99201-99205, 99211-99215, 99241-<br>99245, 99304-99310, 99315, 99316,<br>99318, 99324-99328, 99334-99337,<br>99341-99345, 99347-99350, 99381-<br>99387, 99391-99397, 99401-99404,<br>99411, 99412, 99420, 99429;<br><b>HCPCS:</b> G0402, G0438, G0439,<br>G0463, T1015, S0620, S0621; <b>ICD-</b><br><b>10</b> : Z00.00, Z00.01, Z00.121,<br>Z00.129, Z00.3, Z00.5, Z00.8, Z02.0-<br>Z02.6, Z02.71, Z02.79, Z02.81-<br>Z02.83, Z02.89, Z02.9, Z76.1, Z76.2 |                          |  |
| ADV                            | Annual Dental Visit   | MHT                      | Percentage of members 2-20 years of<br>age who had a dental visit during the<br>measurement year  | Annual dental visit with <b>a dental</b><br><b>provider</b>  |                          |  |

| Code  | Measure Title  | Product Line             | Description  | Coding<br>(Identifying Numerator)   | Category II<br>Codes-CPT   |  |  |
|-------|--|--------------------------|--|---|--|--|--|
| OVERU | OVERUSE/APPROPRIATENESS, CONT.   |                          |  |   |  |  |  |
| IET   |  | Initiation and           |  | Percentage of members 13 years of age and older with a new diagnosis of   | Initiation of AOD treatment: An<br>inpatient admission, outpatient<br>visit, intensive outpatient<br>encounter or partial hospitalization<br>within 14 days of diagnosis -or-<br>CPT: 98966-98968, 99441-99443 |  |  |
|       | Engagement of<br>Alcohol and Other<br>Drug Dependence<br>Treatment                             | ALL                      | alcohol or drug dependence during<br>the measurement year who had<br>initation and engagement of<br>treatment visits within 14 and 34 days<br>of diagnosis.  | <b>Engagement of AOD treatment:</b><br>Initiation of AOD treatment and<br>two or more inpatient admissions,<br>outpatient visits, intensive<br>outpatient encounters or partial<br>hospitalizations with any AOD<br>diagnosis within 34 days after date<br>of the initiation encounter <b>-or- CPT:</b><br>98966-98968, 99441-99443 |  |  |  |
| PPC   | Prenatal and   | Commercial               | Percentage of deliveries with  | <b>Timeliness of prenatal care:</b> a prenatal visit in the first trimester or within 42 days of enrollment   |  |  |  |
|       | Postpartum Care  | & Medicaid               | care   | <b>Postpartum care:</b> a postpartum visit between 7 and 84 days after delivery   |  |  |  |
| APP   | Use of First-Line<br>Psychosocial<br>Care for Children<br>and Adolescents<br>on Antipsychotics | Commercial<br>& Medicaid | Percentage of children and<br>adolescents 1-17 years of age who<br>had a new prescription for an<br>antipsychotic medication and had<br>documentation of psychosocial care<br>as first-line treatment during the<br>measurement year | <b>CPT:</b> 90832-90834, 90836-90840,<br>90845-90847, 90849, 90853, 90875,<br>90875, 90880; <b>HCPCS:</b> G0176,<br>G0177, G0409-G0411, H0004,<br>H0035-H0040, H2000, H2001,<br>H2011-H2014, H2017-H2020, S0201,<br>S9480, S9484, S9485   |  |  |  |

| Code          | Measure Title  | Product Line             | Description  | Coding<br>(Identifying Numerator)  | Category II<br>Codes-CPT |  |  |
|---------------|--|--------------------------|--|--|--------------------------|--|--|
| USE OF        | USE OF SERVICES  |                          |  |  |                          |  |  |
| W34           | Well-Child Visits in<br>the Third, Fourth,<br>Fifth and Sixth<br>Years of Life | Commercial<br>& Medicaid | Percentage of members 3-6 years of<br>age who received one or more well-<br>child visits with PCP during the<br>measurement year               | <b>CPT:</b> 99381-99385, 99391-99395,<br>99461; <b>HCPCS:</b> G0438, G0439;<br><b>ICD-10:</b> Z00.00, Z00.01, Z00.110,<br>Z00.111, Z00.121, Z00.129, Z00.5,<br>Z00.8, Z02.0-ZZ02.6, Z02.71, Z02.82,<br>Z76.1, Z76.2  |                          |  |  |
| AWC           | Adolescent Well-<br>Care Visits  | Commercial<br>& Medicaid | Percentage of members 12-21 years of<br>age who received one or more well-<br>child visit with PCP or OB/GYN during the<br>measurement year    | <b>CPT:</b> 99381-99385, 99391-99395,<br>99461; <b>HCPCS:</b> G0438, G0439;<br><b>ICD-10</b> : Z00.00, Z00.01, Z00.110,<br>Z00.111, Z00.121, Z00.129, Z00.5,<br>Z00.8, Z02.0-ZZ02.6, Z02.71, Z02.82,<br>Z76.1, Z76.2 |                          |  |  |
| PCR           | Plan All Cause<br>Readmissions   | ALL                      | Members 18 years of age and older<br>who had an acute discharge with an<br>unplanned readmission within 30 days<br>during the measurement year |  |                          |  |  |
| <b>RISK A</b> | DJUSTED UTILIZATION  |                          |  |  |                          |  |  |
| AHU           | Acute Hospital<br>Utilization  | Commercial<br>& Medicare | Members 18 and older, the acute<br>impatient and observation stay<br>discharges during the measurement<br>year                                 |  |                          |  |  |
| EDU           | Emergency<br>Department<br>Utilization   | Commercial<br>& Medicare | Members 18 years and older,<br>emergency department visits during the<br>measurement year  |  |                          |  |  |

| Code  | Measure Title  | Product Line             | Description  | Coding<br>(Identifying Numerator)  | Category II<br>Codes-CPT |  |  |
|-------|--|--------------------------|--|--|--------------------------|--|--|
| MEASU | MEASURES COLLECTED THROUGH CAHPS HEALTH PLAN SURVEY      |                          |  |  |                          |  |  |
| FVA   | Flu Shots for Adults                                     | Commercial<br>& Medicaid | Percentage of members 18-64 years<br>who received a flu shot in the last year                                  |  |                          |  |  |
| FVO   | Flu Shots in Older<br>Adults                             | Medicare                 | Percentage of members 65 years and<br>older who received a flu shot in the last<br>year                        |  |                          |  |  |
| FRM   | Fall Risk<br>Management                                  | Medicare                 | Assesses different facets of fall risk<br>management   | Discussing and Managing Fall<br>Risk   |                          |  |  |
| MUI   | Management<br>Urinary<br>Incontinence in<br>Older Adults | Medicare                 | Assesses the management of urinary incontinence in older adults  | Discussing Urinary Incontinence<br>and Treatment   |                          |  |  |
| οτο   | Osteoporosis<br>Testing in Women                         | Medicare                 | Percentage of women 65 years and older who received a bone density test  | Reported ever having received<br>a bone density test to check for<br>osteoporosis  |                          |  |  |
| ΡΑΟ   | Physical Activity in<br>Older Adults                     | Medicare                 | Assesses different facets of promoting physical activity in older adults                                       | Discussing and Advising Physical<br>Activity   |                          |  |  |
| PNU   | Pneumonia<br>Vaccine Status                              | Medicare                 | Percentage of members 65 years and<br>older who received a pneumococcal<br>vaccine during the measurement year |  |                          |  |  |
| MSC   | Medical Assistance<br>with Smoking<br>Cessation          | ALL                      | Assesses different facets of providing<br>medical assistance with smoking and<br>tobacco cessation             | Advising Smokers and Tobacco<br>Users to Quit, Discussing<br>Cessation Medications, and<br>Discussing Cessation Strategies |                          |  |  |