

Air Pollution and Children's Health

Joel Schwartz, PhD

ABSTRACT. Children's exposure to air pollution is a special concern because their immune system and lungs are not fully developed when exposure begins, raising the possibility of different responses than seen in adults. In addition, children spend more time outside, where the concentrations of pollution from traffic, powerplants, and other combustion sources are generally higher. Although air pollution has long been thought to exacerbate minor acute illnesses, recent studies have suggested that air pollution, particularly traffic-related pollution, is associated with infant mortality and the development of asthma and atopy. Other studies have associated particulate air pollution with acute bronchitis in children and demonstrated that rates of bronchitis and chronic cough declined in areas where particle concentrations have fallen. More mixed results have been reported for lung function. Overall, evidence for effects of air pollution on children have been growing, and effects are seen at concentrations that are common today. Although many of these associations seem likely to be causal, others require and warrant additional investigation. *Pediatrics* 2004;113:1037-1043; *asthma, particles, ozone, lung reaction.*

ABBREVIATIONS. PM₁₀, particles with aerodynamic diameter less than 10 μ m; NO, nitric oxide; CI, confidence interval.

The health effects of air pollution exposure have become an area of increasing focus in the past 30 years. A growing body of evidence has demonstrated that there are serious health consequences to community air pollution and that these consequences are not spread equally among the population. As an example of this differential susceptibility, recent studies have indicated that people with type 2 diabetes are at higher risk for cardiovascular effects of airborne particles.^{1,2} Similarly, children have been shown to be at particular risk for other effects of air pollution, as detailed below.

This article cannot be a comprehensive review of the literature, because recent reviews of airborne particles and ozone alone have hundreds of pages summarizing the literature. Rather, I cover the major health effects in children that have been linked to air pollution, cite some key papers, and discuss the strength of the evidence. I particularly highlight ar-

reas where it seems that differences between adults and children, particularly in the development of the respiratory and immune system, suggest different impacts of exposure for children.

BACKGROUND

The lung is not well formed at birth, and development of full functionality does not occur until approximately 6 years of age. During early childhood, the bronchial tree is still developing. For example, the number of alveoli in the human lung increases from 24 million at birth to 257 million at age 4,³ and the lung epithelium is not fully developed. This results in greater permeability of the epithelial layer in young children. Children also have a larger lung surface area per kilogram of body weight than adults and, under normal breathing, breathe 50% more air per kilogram of body weight than adults. This process of early growth and development, the outcome of which is important for the future health of the child, suggests that there is a critical exposure time when air pollution may have lasting effects on respiratory health.

At the same time the child's lung is developing, the child's immune system, immature at birth, is also beginning to develop. Much recent attention in asthma research has been focused on this development, in particular factors that influence the development of TH-2 (humoral immunity dominant) versus TH-1 (cellular immunity dominant) phenotypes.⁴

Another major factor that influences the relative impact of air pollution on children versus adults is exposure. Children spend more time outdoors than adults, particularly in the summer and in the late afternoon.⁵ Some of that time is spent in activities that increase ventilation rates. This can increase the exposure to air pollutants compared with adults, as indoor concentrations of air pollutants of outdoor origin are usually lower.

PRE- AND PERINATAL EFFECTS OF AIR POLLUTION

Although historically air pollution has been thought of as a respiratory toxicant, recent evidence has broadened our understanding of its full range of effects. In adults, changes in cardiovascular risk factors such as C-reactive protein⁶ and autonomic control of the heart⁷ have led the way in broadening our understanding of the range of toxicity. With children, perhaps the most unexpected results have been a range of recent papers reporting that prenatal exposure of populations to prevailing levels of air pollution is associated with early fetal loss,⁸ preterm

From the Departments of Environmental Health and Epidemiology, Harvard School of Public Health, and Channing Laboratory, Brigham and Women's Hospital, Department of Medicine, Harvard Medical School, Boston, Massachusetts.

Received for publication Oct 7, 2003; accepted Oct 20, 2003.

Reprint requests to (J.S.) Environmental Epidemiology Program, Harvard School of Public Health, 401 Park Dr, Boston, MA 02215. E-mail: jschwartz@hsph.harvard.edu

PEDIATRICS (ISSN 0031 4005). Copyright © 2004 by the American Academy of Pediatrics.

delivery,^{9–11} and lower birth weight.^{12–18} These associations may or may not be causal but clearly warrant additional study. The later Bobak study¹⁸ is notable in that it was nested within a birth cohort study, allowing good control for social and other factors that may confound the association. Because birth certificates in most areas have extensive information on maternal medical conditions that may affect the pregnancy, as well as maternal age, education, and smoking, all of these studies are generally well controlled. Although relatively recent, there is now considerable evidence that maternal exposure to air pollution during pregnancy is associated with adverse birth outcomes. Moreover, particulate air pollution from combustion sources shares many characteristics with sidestream tobacco smoke, which is rich in particles and polycyclic aromatic hydrocarbons. A recent review by Windham et al¹⁹ found that environmental tobacco smoke was associated with low birth weight. This provides support for the plausibility of the reported association.

The mechanisms involved are as yet unknown but may include inflammatory processes and oxidative stress, which have been linked to air pollution. For example, Salvi et al²⁰ reported that human volunteers who were exposed to diesel particles for 1 hour had elevated levels of peripheral white cells, as well as increased vascular cellular adhesion molecule-1 and intercellular adhesion molecule-1 in the lung epithelium. As noted before, C-reactive protein, an acute-phase inflammatory marker, was associated with air pollution exposure in adults. Ozone is a highly reactive gas, associated with oxidative stress in many studies.^{21–24}

Additional support is provided by some animal studies, which provide some ideas about mechanisms. Although these tend to be at high doses, they can help to supplement the human data. Recently, Saldiva and co-workers^{25,26} reported lung inflammation associated with particle and particle component exposure in rats. Carbon monoxide exposure has been associated with fetal toxicity, including intra-uterine growth retardation in the rat.²⁷ Ozone exposure has also been shown to be fetotoxic in an animal model.²⁸

Perhaps the most serious thing that can be done to a child's life is to end it. Recently, a number of studies have reported that air pollution is associated with precisely that. In thinking about air pollution and death, one is inevitably led to the great air pollution episode of December 1–5, 1952, in London. A low-level thermal inversion that trapped coal smoke in the Thames valley, coupled with a stationary front that dropped wind speed to 0, resulted in a rapid buildup of pollution to extremely high levels. Approximately 4000 excess deaths occurred in London in that week,²⁹ and elevated death rates continued for weeks afterward,³⁰ indicating that there were delayed as well as prompt effects. Although most of the deaths were in adults, infant mortality was doubled during that period.³¹

This episode is important because it establishes causality. The influenza epidemic did not arrive in England until >1 month after the episode, and in

other towns in England, where the weather was as cold or colder but no inversion occurred, no increase in deaths was observed. Furthermore, the death rate increased rapidly in phase with the pollution and began to come down when the pollution came down. Hence, it is clear that at very high levels, air pollution can produce a substantial increase in deaths of children.

More recently, Woodruff et al³² examined infant deaths in the United States and levels of inhalable particles (PM₁₀) in the air. They excluded infant deaths in the first month after birth as likely to reflect complications of pregnancy and delivery and found that PM₁₀ was associated with higher death rates in the next 11 months of life. This excess risk seemed to be principally from respiratory illness, although sudden infant death syndrome deaths were also elevated.

Bobak and Leon³³ recently also examined the cross-sectional association between air pollution and infant mortality rates across towns in the Czech Republic. A significant association was seen between infant death rates and particle and SO₂ concentration. Other studies have examined day-to-day changes in air pollution and day-to-day changes in infant deaths. Saldiva et al³⁴ reported that infant death from respiratory disease was associated with air pollution, particularly from traffic. Loomis et al³⁵ similarly found respiratory deaths in infants associated with air pollution.

ACUTE EFFECTS OF AIR POLLUTION EXPOSURE

Exposure Issues

As noted above, children's exposure can be different from adults given the same outdoor concentrations. This is particularly important for exposure to ozone. Ozone is a highly reactive gas, producing oxidative damage in the lung. Because of that high reactivity, its half-life in indoor air is only 7 to 10 minutes.³⁶ Consequently, ozone levels are very low indoors.^{5,37} This is particularly true for locations with low air exchange rates, such as air-conditioned homes and workspaces. Ozone also has a distinct temporal pattern. Because it is not directly emitted from polluting sources but produced by photochemical reactions in the atmosphere, it shows strong seasonal and diurnal variations. It is high in the summer and the afternoon and low in the night, early morning, and winter. Children tend to be outdoors in the afternoon and in the summer, which results in much higher exposure for children than adults, who are protected by their indoor environment.

In contrast, fine combustion particles, usually indexed by PM_{2.5} (particles <2.5 μm in aerodynamic diameter) penetrate indoors and are not chemically quenched like ozone (or SO₂). Recent studies of the association between personal exposure to particles of outdoor origins and outdoor concentrations show that the personal exposures are much more tightly linked than for ozone,³⁸ although they do vary with air exchange rates of the buildings in which the person spends time. Hence children's exposure to

PM_{2.5} is enhanced by their greater outdoor activity for this pollutant but to a lesser extent than for ozone.

Health Effects

There is a large body of literature associating short-term changes in air pollution with short-term changes in pulmonary health of children, often focused on individuals with asthma. Of particular interest are a series of summer camp studies^{39–41}. These were innovative because the living conditions of the children in the camp meant that they were exposed all the time. For most of the day, they were outdoors, and their indoor quarters had such high air infiltration rates that indoor concentrations of outdoor pollutants were almost certainly similar to the outdoor levels. In these studies, lung function declined during air pollution episodes, which were combinations of ozone and sulfate particulates, some of which may have been acidic.

Another set of studies examined wintertime episodes. A study in Steubenville, Ohio, repeated measurements of pulmonary function in schoolchildren before, during, and after an episode of high-particulate air pollution.⁴² Lung function declined during the episode. A similar study was performed in the Netherlands.⁴³

These studies were followed up by a different study design that made it possible to collect large amounts of data relatively inexpensively. A panel of children would be recruited and asked to perform daily peak flow tests and usually to answer questions on symptom prevalence (wheezing, coughing, etc) for a period of several months. These measurements were then correlated with air pollution. Often but not always, these studies would be focused on children with asthma. In general, significant associations have been reported with PM₁₀,^{44–48} although not in every study.⁴⁹ Other summer time studies have reported associations with ozone,^{50–52} including interactions with aeroallergen exposure.⁵³

Of particular interest are 2 studies from the Netherlands that addressed the question of susceptibility. Van der Zee et al⁵⁴ looked at wintertime air pollution in panels of 7- to 11-year-old children. A stronger association was found between particle pollution and peak flow decrements in children with asthma symptoms, particularly those on regular medication, than with nonsymptomatic children. There was also a significant effect on bronchodilator use in the symptomatic children. A second analysis looked at more objective measures than reports of chronic respiratory symptoms as effect modifiers. Children were stratified into those with measured bronchial hyperresponsiveness and elevated immunoglobulin E and those without.⁵⁵ The association between decrements in peak flow and air pollution was primarily in the former group.

A related approach is to use administrative data to look at more serious outcomes that require physician contact. For example, Pope⁵⁶ examined hospital admissions of children in the Utah valley during 3 consecutive winters. These winters were before, during, and after a strike that closed down a steel mill in the valley that was the largest source of wintertime

air pollution. There was a >50% drop in admissions of children for asthma and for pneumonia during the period that the mill was closed and when the pollution was lower. In the following year, admissions went back to previous levels. In a neighboring valley, there was no drop in pollution or admissions in the middle winter. This is as close to a clinical trial as can be found in air pollution epidemiology, and the conclusions are striking. Air pollution is related to serious asthma exacerbation and to pneumonia exacerbation. Other studies have looked at day-to-day fluctuations in hospital admissions and day-to-day changes in air pollution and reported associations with childhood hospital admissions in Ontario,^{57,58} Seattle,^{59,60} and elsewhere.^{61,62}

A different approach is to look at physician visits. Such data are hard to obtain systematically for large populations in the United States but are more readily available in Europe. Medina et al⁶³ looked at emergency house calls by physicians in Paris and found that visits for asthma were associated with particulate air pollution and ozone and that the association was stronger for children.

What evidence is there that these associations are plausible? An important study by Zelikoff et al⁶⁴ showed that exposure to urban particles exacerbated pneumonia in an animal model, supporting the results of the epidemiologic studies in Utah and elsewhere. Other studies have shown ozone to be associated with altered macrophage function and epithelial injury,⁶⁵ which could plausibly modify infectivity.

Other evidence points to a role for pollution in increasing lung inflammation in children, particularly those with asthma. For example, Fischer et al⁶⁶ examined a cohort of 68 children (aged 10–11) with 7 weekly measurements of exhaled nitric oxide (NO) and found that increases in several air pollutants were associated with increased levels of exhaled NO, a good marker of lung inflammation in individuals with asthma.^{67,68} Giroux et al⁶⁹ contrasted exhaled NO in children who had asthma and lived in urban areas with others who stayed in a national park in the mountains, or with children without asthma in the same city. The exhaled NO concentrations in the urban children with asthma were more than double those in the children with asthma in the national park, and their was no difference in exhaled NO between children with asthma in the park and healthy children in the city.

Finally, we have excellent evidence that changing pollution in the short term produces immediate reductions in asthma exacerbations. In addition to the Utah study cited above, a more recent study looked at asthma hospital visits in Atlanta around the period of the Olympics, when traffic was limited and air pollution was reduced. A noticeable reduction in asthma emergency visits occurred during that period of short-term traffic control.⁷⁰

EFFECTS OF LONG-TERM EXPOSURE TO AIR POLLUTION

Although the role of air pollution in exacerbating existing illness has been well established, recent ev-

idence has implicated pollution exposure with the development of chronic disease or impairments. Evidence has been accumulating for a while about effects on lung function and bronchitic symptoms. More recently, studies have begun to implicate air pollution, particularly from traffic, with the pathogenesis of asthma.

In the late 1980s, Schwartz⁷¹ examined the association between long-term exposure of children to air pollution and pulmonary function in the Second National Health and Nutrition Examination Survey. He found significant decrements in lung function associated with exposure.

Jedrychowski et al⁷² also reported that air pollution was associated with lower levels of lung function growth in children in Poland. Horak et al⁷³ made repeated measurements of spirometry during a 3-year period in Austrian schoolchildren and found that after adjustment for covariates, including initial lung function, lung function growth rates were associated with PM₁₀ exposure. An increase of 10 µg/m³ in PM₁₀ exposure was associated with a decrease in growth of forced expiratory volume in 1 second of 84 mL/year.

Other studies have implicated ozone exposure during childhood with reductions in lung function. For example, Künzli et al⁷⁴ collected residential address histories for freshmen at the University of California at Berkeley and matched them to monitors near their homes. Cumulative ozone exposure was associated with a significant decrement in forced expiratory volume in 1 second. A similar result was found for freshmen at Yale.⁷⁵

Dockery et al⁷⁶ reported that chronic bronchitis and chest illness in children were associated with exposure to particulate air pollution. This study compared covariate adjusted rates across 6 communities in the eastern United States with varying levels of pollution. No association was seen with asthma or wheezing. Subsequent studies in the United States⁷⁷ and elsewhere confirmed that particulate exposure was associated with higher rates of chronic cough and bronchitis symptoms in children and the lack of association with wheezing and asthma. A similar large study (*n* = 4470) comparing schoolchildren in 10 communities in Switzerland reported an adjusted odds ratio for bronchitis of 2.88 (95% confidence interval [CI]: 1.69–4.89) for PM₁₀ exposure between the most and least polluted community.⁷⁸ A study of 3676 children across 12 Southern California communities reported that bronchitis was associated with PM₁₀ but only among children with asthma.⁷⁷ The largest study examined 13 369 children in 24 communities in the United States and Canada.⁷⁹ Again, particulate air pollution was associated with bronchitis episodes across these communities.

A recent study that looked at eastern Germany, where there has been a reduction in pollution since the reunification, shows that this reduction has been associated with reductions in the rates of chronic cough and bronchitis symptoms in a new cohort of children.⁸⁰ This demonstrates not merely an association but that an intervention produces improvements in health. A similar dramatic effect of intervention

was seen in a study by Avol et al.⁸¹ Using the Southern California cohort study mentioned above, they identified 110 children who moved from the study area and followed them up in their new home with pulmonary function testing identical to that in the main cohort. Subjects who moved to locations with higher PM₁₀ concentrations showed lower rates of annual growth in lung function, and subjects who moved to locations with lower PM₁₀ concentrations than they had left showed higher rates of growth in lung function. This effect was increased in subjects who lived in the new location for at least 3 years.

Of considerable interest are recent studies that have called into question the previous results indicating that long-term air pollution exposure (mostly to particles) was associated with bronchitis symptoms but not asthma. These studies all used central monitoring locations in each community to assess long-term exposure in those communities. Although these monitoring stations are reasonable surrogates for long-term exposure to pollutants that are relatively homogeneously distributed across the community, that is not true for all pollutants. In particular, traffic pollutants show strong gradients. Exposure to diesel exhaust varies greatly with distance from major roadways within a community.^{82,83} The new studies have used measurements or models of this microlevel spatial variability in exposure within community and returned to the question of whether air pollution exposure is associated with the development of asthma.

Studnicka et al⁸⁴ examined 8 small, rural communities with no industrial sources of pollution. NO₂ was measured in each community and taken as a measure of exposure to traffic pollution. In areas without heavy industry, almost all NO₂ is attributable to traffic. Although both gasoline and diesel engines produce NO₂, diesel engines produce much more, so this surrogate is weighted toward diesel exposure. A strong association between asthma prevalence and NO₂ levels was found, with odds ratios reaching 5.81 (95% CI: 1.27–26.5), contrasting the highest and lowest exposures. Kramer et al⁸⁵ examined 317 children in 3 German communities. NO₂ measurements were made outside the homes of each of the children, and personal NO₂ measurements were collected for each child. The personal NO₂ measurements reflect exposure to both outdoor NO₂ and indoor sources (eg, gas stoves). The NO₂ outside the home reflected exposure to NO₂ from outdoor sources only and therefore was a good surrogate for exposure to traffic pollution. The NO₂ measurements outside each child's home were significant predictors of hay fever; symptoms of allergic rhinitis; wheezing; and sensitization against pollen, house dust mites, or cats. The personal NO₂ measurements, which were strongly influenced by indoor sources, were not associated with these outcomes. This indicates that traffic pollution but probably not the NO₂ from traffic is associated with atopy and wheezing. If NO₂ per se is not the relevant exposure, than diesel particles or some component of those particles, such as polycyclic aromatic hydrocarbons, may be the most important etiologic compo-

ment. In the Netherlands, residence on a high-traffic street was associated with a >2-fold increase in the risk of wheezing after control for confounders.⁸⁶

Even more recently, Lin et al⁸⁷ geocoded the residential addresses of children who were admitted to the hospital in Erie County, New York (excluding Buffalo) for asthma, and age-matched controls who were admitted for nonrespiratory conditions. These were linked to Department of Transportation data on vehicle miles traveled on their street. The odds of asthma (adjusted for poverty level) for living within 200 m of a street with the highest tertile of traffic density was 1.93 (95% CI: 1.13–3.29), and the children with asthma were more likely to have truck traffic on their street. Another recent report analyzed data from 2 birth cohorts totaling 1756 children in Munich.⁸⁸ Geographic Information System modeling was used to estimate the concentrations of traffic-related particles and NO₂ outside the birth addresses of all of the children. These pollutants were associated with dry cough at night in the first year of life. Another case-control study of 6147 children in Nottingham, England, found increased risk of wheeze associated with living within 90 m of a roadway.⁸⁹ Although some studies showed no increased risk,⁹⁰ the overwhelming weight of the recent evidence suggests that traffic pollution is associated with the risk of developing asthma.

CONCLUSIONS

Air pollution is not the leading cause of death or morbidity in children in the developed world. However, there is increasingly strong evidence that air pollution is associated with nontrivial increases in the risk of death and chronic disease in children, worse pregnancy outcomes, and exacerbation of illnesses. It is less clear which pollutants are most responsible, but particles and ozone have the strongest associations. For the incidence of asthma, traffic pollution, particularly from trucks, seems to be the key player.

What is important to realize is that this is an easily modifiable risk. Sulfate particles, a major fraction of the particle burden in the air in urban areas, can be easily removed using scrubbers on powerplants (their largest source) at a cost that is <1% of the current price of electricity. NO_x reduction, a major component of an ozone reduction strategy, can also be retrofitted onto powerplants. In Europe, catalytic converters on cars can be brought up to US standards. Traffic particles, NO_x, and so forth are dominated by diesel engines. Trap oxidizers and catalysts can reduce these emissions by up to 90%. Such devices have been on gasoline-powered vehicles for decades without ending industrial civilization as we know it. For many of these control strategies, it does not matter that we are not sure which component of the pollution mix is principally responsible. Oxidative catalysts reduce carbon soot, polycyclic aromatic hydrocarbons, CO, and so forth. Given the amount of money that we spend on the treatment of asthma and the difficulty that we have in reducing allergen exposures, such straightforward approaches need serious attention.

REFERENCES

1. Zanobetti A, Schwartz J. Are diabetics more susceptible to the health effects of airborne particles? *Am J Respir Crit Care Med.* 2001;164:831–833
2. Bateson TF, Schwartz J. Who is sensitive to the effects of particles on mortality? A case-crossover analysis of individual characteristics as effect modifiers. *Epidemiology.* 2004, in press
3. Dunnill MS. Thorax 1962;17:329, cited in: Altman PL, Dittmer DS. *Respiration and Circulation.* Bethesda, MD: Federation of American Societies for Experimental Biology; 1971
4. Holt PG. Programming for responsiveness to environmental antigens that trigger allergic respiratory disease in adulthood is initiated in the perinatal period. *Environ Health Perspect.* 1998;106(suppl 3):795–800
5. US Environmental Protection Agency, National Center for Exposure Assessment. *Exposure Factors Handbook.* Washington, DC: Environmental Protection Agency; 1997 (EPA/600/P95/002Fa)
6. Peters A, Frohlich M, Doring A, et al. Particulate air pollution is associated with an acute phase response in men; results from the MONICA-Augsburg Study. *Eur Heart J.* 2001;22:1198–1204
7. Gold DR, Litonjua A, Schwartz J, et al. Ambient pollution and heart rate variability. *Circulation.* 2000;101:1267–1273
8. Pereira LA, Loomis D, Conceicao GM, et al. Association between air pollution and intrauterine mortality in Sao Paulo, Brazil. *Environ Health Perspect.* 1998;106:325–329
9. Xu X, Ding H, Wang X. Acute effects of total suspended particles and sulfur dioxides on preterm delivery: a community-based cohort study. *Arch Environ Health.* 1995;50:407–415
10. Ritz B, Yu F, Chapa G, Fruin S. Effect of air pollution on preterm birth among children born in southern California between 1989 and 1993. *Epidemiology.* 2000;11:502–511
11. Lin MC, Chiu HF, Yu HS, et al. Increased risk of preterm delivery in areas with air pollution from a petroleum refinery plant in Taiwan. *J Toxicol Environ Health A.* 2001;64:37–44
12. Wang X, Ding H, Ryan L, Xu X. Association between air pollution and low birth weight: a community based study. *Environ Health Perspect.* 1997;105:514–520
13. Ritz B, Yu F. The effect of ambient carbon monoxide on low birth weight among children born in southern California between 1989 and 1993. *Environ Health Perspect.* 1999;107:17–25
14. Dejmeck J, Selevan SG, Benes I, Solansky I, Sram RJ. Fetal growth and maternal exposure to particulate matter during pregnancy. *Environ Health Perspect.* 1999;107:475–480
15. Bobak M, Leon DA. Pregnancy outcomes and outdoor air pollution: an ecological study in districts of the Czech republic 1986–1988. *Occup Environ Med.* 1999;56:539–543
16. Ha EH, Hong YC, Lee BE, Woo BH, Schwartz J, Christiani DC. Is air pollution a risk factor for low birth weight in Seoul? *Epidemiology.* 2001;12:643–648
17. Bobak M. Outdoor air pollution, low birth weight, and prematurity. *Environ Health Perspect.* 2000;108:539–543
18. Bobak M, Richards M, Wadsworth M. Air pollution and birth weight in Britain in 1946. *Epidemiology.* 2001;12:358–359
19. Windham GC, Eaton A, Hopkins B. Evidence for an association between environmental tobacco smoke exposure and birthweight: a meta-analysis and new data. *Paediatr Perinat Epidemiol.* 1999;13:35–57
20. Salvi S, Blomberg A, Rudell B, et al. Acute inflammatory responses in the airways and peripheral blood after short-term exposure to diesel exhaust in healthy human volunteers. *Am J Respir Crit Care Med.* 1999; 159:702–709
21. Pryor WA, Church DF. The reaction of ozone with unsaturated fatty acids: aldehydes and hydrogen peroxide as mediators of ozone toxicity. In: Davies KJA, ed. *Oxidative Damage and Repair: Chemical, Biological and Medical Aspects.* New York, NY: Pergamon Press; 1991:496–504
22. Koren HS, Devlin RB, Graham De, et al. Ozone induced inflammation in lower airways of human subjects. *Am Rev Respir Dis.* 1989;139:407–415
23. Mudway IS, Krishna MT, Frew AJ, et al. Compromised concentrations of ascorbate in fluid lining the respiratory tract in human subjects after exposure to ozone. *Occup Environ Med.* 1999;56:473–481
24. Saintot M, Bernard N, Astre C, Gerber M. Ozone exposure and blood antioxidants: a study in a periurban area in Southern France. *Arch Environ Health.* 1999;54:34–39
25. Saldiva PH, Clarke RW, Coull BA, et al. Lung inflammation induced by concentrated ambient air particles is related to particle composition. *Am J Respir Crit Care Med.* 2002;165:1610–1617
26. Rice TM, Clarke RW, Godleski JJ, et al. Differential ability of transition metals to induce pulmonary inflammation. *Toxicol Appl Pharmacol.* 2001; 177:46–53

27. Garvey DJ, Longo LD. Chronic low level maternal carbon monoxide exposure and fetal growth and development. *Biol Reprod.* 1978;19:8–14
28. Kavlock R, Daston G, Grabowski CT. Studies on the developmental toxicity of ozone. I. Prenatal effects. *Toxicol Appl Pharmacol.* 1979; 48(suppl):19–28
29. Mortality and morbidity during the London fog of December 1952. Her Majesty's Stationary Office, London. Report No. 95 on Public Health and Medical Subjects. London, UK: Her Majesty's Public Health Service; 1954
30. Anderson HR. Health effects of air pollution episodes. In: Holgate ST, Samet JM, Koren HS, Maynard RL, eds. *Air Pollution and Health.* London, UK: Academic Press; 1999:461–484
31. Schwartz J. What are people dying of on high air pollution days? *Environ Res.* 1994;64:26–35
32. Woodruff TJ, Grillo J, Schoendorf C. The relationship between selected causes of postneonatal infant mortality and particulate air pollution in the United States. *Environ Health Perspect.* 1997;105:608–612
33. Bobak M, Leon DA. Air pollution and infant mortality in the Czech Republic, 1986–88. *Lancet.* 1992;340:1010–1014
34. Saldiva PHN, Lichtenfels AJFC, Paiva PSO, et al. Association between air pollution and mortality due to respiratory disease in children in Sao Paulo, Brazil. *Environ Res.* 1994;65:218–225
35. Loomis D, Castillejos M, Gold DR, McDonnell W, Borja-Aburto VH. Air pollution and infant mortality in Mexico City. *Epidemiology.* 1999;10: 118–123
36. Weschler CJ. Ozone in indoor environments: concentration and chemistry. *Indoor Air.* 2000;10:269–288
37. Lee K, Xue J, Geyh AS, et al. Nitrous acid, nitrogen dioxide, and ozone concentrations in residential environments. *Environ Health Perspect.* 2002;110:145–150
38. Sarnat JA, Schwartz J, Catalano PJ, Suh HH. Confounder or surrogate: the role of gaseous pollutants in particulate matter epidemiology. *Environ Health Perspect.* 2001;109:1053–1061
39. Spektor DM, Mao J, He D, Thurston GD, Hayes C, Lippmann M. Effects of single and multi day ozone exposures on respiratory function in active normal children. *Environ Res.* 1991;55:107–122
40. Kinney PL, Ware JH, Spengler JD, Dockery DW, Speizer FE, Ferris BG Jr. Short-term pulmonary function change in association with ozone levels. *Am Rev Respir Dis.* 1989;139:56–61
41. Berry M, Lioy PJ, Gelperin K, Buckler G, Klotz J. Accumulated exposure to ozone and measurement of health effects in children and counselors at two summer camps. *Environ Res.* 1991;54:135–150
42. Dockery DW, Ware JH, Ferris BG, Speizer FE, Cook NR, Herman SM. Change in pulmonary function associated with air pollution episodes. *J Air Poll Control Assoc.* 1982;32:937–942
43. Dassen W, Brunekreef B, Hoek G, et al. Decline in children's pulmonary function during an air pollution episode. *J Air Poll Control Assoc.* 1986; 36:1123–1127
44. Romieu I, Meneses F, Ruiz S, et al. Effects of air pollution on the respiratory health of asthmatic children living in Mexico City. *Am J Respir Crit Care Med.* 1996;154:300–307
45. Ostro B, Lipsett M, Mann J, Braxton-Owens H, White M. Air pollution exacerbation of asthma in African-American children in Los Angeles. *Epidemiology.* 12:200–208
46. Pope CA, Dockery DW. Acute health effects of PM₁₀ pollution on symptomatic and asymptomatic children. *Am Rev Respir Dis.* 145: 1123–1128
47. Braun-Fahrlander C, Ackerman-Lieblich U, Schwartz J, Grehm HP, Rutishauser M, Wanner HU. Air pollution and respiratory symptoms in pre-school children. *Am Rev Respir Dis.* 1992;145:42–47
48. Schwartz J, Dockery DW, Neas LM, et al. Acute effects of summer air pollution on respiratory symptom reporting in children. *Am J Respir Crit Care Med.* 1994;150:1234–1242
49. Roemer W, Clench-Aas J, Englert N, et al. Inhomogeneity in response to air pollution in European children (PEACE project). *Occup Environ Med.* 1999;56:86–92
50. Kinney PL, Lippmann M. Respiratory effects of seasonal exposures to ozone and particles. *Arch Environ Health.* 2000;55:210–216
51. Jalaludin BB, Chey T, O'Toole BI, Smith WT, Capon AG, Leeder SR. Acute effects of low levels of ambient ozone on peak expiratory flow rate in a cohort of Australian children. *Int J Epidemiol.* 2000;29:549–557
52. Gold DR, Damokosh AI, Pope CA 3rd, et al. Particulate and ozone pollutant effects on the respiratory function of children in southwest Mexico City. *Epidemiology.* 1999;10:8–16
53. Higgins BG, Francis HC, Yates C, et al. Environmental exposure to air pollution and allergens and peak flow changes. *Eur Respir J.* 2000;16: 61–66
54. Van der Zee S, Hoek G, Boezen HM, Schouten JP, van Wijnen JH, Brunekreef B. Acute effects of urban air pollution on respiratory health of children with and without chronic respiratory symptoms. *Occup Environ Med.* 1999;12:802–812
55. Boezen HM, van der Zee SC, Postma DS, et al. Effects of ambient air pollution on upper and lower respiratory symptoms and peak expiratory flow in children. *Lancet.* 1999;353:874–878
56. Pope CA III. Respiratory disease associated with community air pollution and a steel mill, Utah valley. *Am J Public Health.* 1989;79:623–628
57. Bates DV, Szto R. The Ontario Air Pollution Study: identification of the causative agent. *Environ Health Perspect.* 1989;79:69–72
58. Burnett RT, Dales RE, Raizenne ME, et al. Effects of low ambient levels of ozone and sulfates on the frequency of respiratory admissions to Ontario hospitals. *Environ Res.* 1994;65:172–194
59. Schwartz J, Koenig J, Slater D, Larson T. Particulate air pollution and hospital emergency visits for asthma in Seattle. *Am Rev Respir Dis.* 1993;147:826–831
60. Norris G, Larson T, Koenig J, Claiborn C, Sheppard L, Finn D. Asthma aggravation, combustion, and stagnant air. *Thorax.* 2000;55:466–470
61. Tenias JM, Ballester F, Rivera ML. Association between hospital emergency visits for asthma and air pollution in Valencia Spain. *Occup Environ Med.* 1998;55:541–547
62. Sunyer J, Spix C, Quénel P, et al. Urban air pollution and emergency admissions for asthma in four European cities: the APHEA Project. *Thorax.* 1997;52:760–765
63. Medina S, Le Tertre A, Quénel P, et al. Air pollution and doctors' house calls: results from the ERPURS system for monitoring the effects of air pollution on public health in Greater Paris, France 1991–1995. *Environ Res.* 1997;75:73–84
64. Zelikoff JT, Nadziejko C, Fang T, Gordon C, Premdass C, Cohen MD. Short term, low-dose inhalation of ambient particulate matter exacerbates ongoing pneumococcal infections in Streptococcus Pneumoniae-infected rats. In: Phalen RF, Bell YM, eds. *Proceedings of the Third Colloquium on Particulate Air Pollution and Human Health.* Irvine, CA: Air Pollution Health Effects Laboratory, University of California; 1999: 8–94–8–101
65. Devlin RB, McDonnell WF, Mann R, et al. Exposure of humans to ambient levels of ozone for 6.6 hours causes cellular and biochemical changes in the lung. *Am J Respir Cell Mol Biol.* 1991;4:72–81
66. Fischer PH, Steerenerg PA, Smelder JD, Van Loveren H, Van Amsterdam JG. Association between exhaled nitric oxide, ambient air pollution, and respiratory health in school children. *Int Arch Occup Environ Health.* 2002;75:348–353
67. Kharitonov SA, Yates D, Springall DR, et al. Exhaled nitric oxide is increased in asthma. *Chest.* 1995;107(3 suppl):156S–157S
68. Massaro AF, Mehta S, Lilly CM, Kobzik L, Reilly JJ, Drazen JM. Elevated nitric oxide concentrations in isolated lower airway gas of asthmatic subjects. *Am J Respir Crit Care Med.* 1996;153:1510–1514
69. Giroux M, Bremont F, Ferrieres J, Dumas JC. Exhaled NO in asthmatic children in unpolluted and urban environments. *Environ Int.* 2001;27: 335–340
70. Friedman MS, Powell KE, Hutwagner L, Graham LM, Teague WG. Impact of changes in transportation and commuting behaviors during the 1996 Summer Olympic Games in Atlanta on air quality and childhood asthma. *JAMA.* 2001;285:897–905
71. Schwartz J. Lung function and chronic exposure to air pollution: a cross-sectional analysis of NHANES II. *Environ Res.* 1989;50:309–321
72. Jedrychowski W, Flak E, Mroz E. The adverse effect of low levels of ambient air pollutants on lung function growth in preadolescent children. *Environ Health Perspect.* 1999;107:669–674
73. Horak F, Studnicka M, Gartner C, et al. Particulate matter and lung function growth in children: a three year followup study in Austrian schoolchildren. *Eur Respir J.* 2002;19:838–845
74. Künzli N, Lurmann F, Segal M, Ngl L, Balmes J, Tager IB. Association between lifetime ambient ozone exposure and pulmonary function in college freshmen: results of a pilot study. *Environ Res.* 1997;72:8–23
75. Galizia A, Kinney PL. Long-term residence in areas of high ozone: associations with respiratory health in a nationwide sample of non-smoking young adults. *Environ Health Perspect.* 1999;107:675–679
76. Dockery DW, Speizer FE, Stram DO, Ware JH, Spengler JD, Ferris BG Jr. Effects of inhaled particles on respiratory health of children. *Am Rev Respir Dis.* 1989;139:587–594
77. McConnell R, Berhane K, Gilliland F, et al. Air pollution and bronchitic symptoms in Southern California children with asthma. *Environ Health Perspect.* 1999;107:757–760
78. Braun-Fahrlander C, Vuille JC, Sennhauser FH, et al. Respiratory health and long term exposure to air pollutants in Swiss schoolchildren. SCARPOL team. *Am J Respir Crit Care Med.* 1997;155:1042–1049
79. Dockery DW, Cunningham J, Damokosh AI, et al. Health effects of acid

- aerosols on North American children: respiratory symptoms. *Environ Health Perspect.* 1996;105:500–505
80. Heinrich J, Hoelscher B, Wichman HE. Decline of ambient air pollution and respiratory symptoms in children. *Am J Respir Crit Care Med.* 2000;161:1930–1936
81. Avol EL, Gauderman WJ, Tan SM, London SJ, Peters JM. Respiratory effects of relocating to areas of differing air pollution levels. *Am J Respir Crit Care Med.* 2001;164:2067–2072
82. Roorda-Knape MC, Janssen NA, de Hartog J, Van Vliet PH, Harssema H, Brunekreef B. Air pollution from traffic in city districts near motorways. *Atmos Environ.* 1998;32:1921–1930
83. Levy JI, Dumyahn T, Spengler JD. Particulate matter and polycyclic aromatic hydrocarbon concentrations in indoor and outdoor microenvironments in Boston, Massachusetts. *J Expo Anal Environ Epidemiol.* 2002;12:104–114
84. Studnicka M, Hackl E, Pischinger J, et al. Traffic related NO₂ and the prevalence of asthma and respiratory symptoms in seven year olds. *Eur Respir J.* 1997;10:2275–2278
85. Kramer U, Koch T, Ranft U, Ring J, Behrendt H. Traffic related air pollution is associated with atopy in children living in urban areas. *Epidemiology.* 2000;11:64–70
86. Oosterlee A, Drijver M, Lebet E, Brunekreef B. Chronic respiratory symptoms in children and adults living along streets with high traffic density. *Occup Environ Med.* 1996;53:241–247
87. Lin S, Munsie JP, Hwang SA, Fitzgerald E, Cayo MR. Childhood asthma hospitalization and residential exposure to state route traffic. *Environ Res.* 2002;88:73–81
88. Gehring U, Cyrys J, Sedlmeir G, et al. Traffic related air pollution and respiratory health during the first 2 yrs of life. *Eur Respir J.* 2002;19:690–698
89. Venn AJ, Lewis SA, Cooper M, Hubbard R, Britton J. Living near a main road and the risk of wheezing illness in children. *Am J Respir Crit Care Med.* 2001;164:2177–2180
90. English P, Neutra R, Scalf R, Sullivan M, Waller L, Zhu L. Examining associations between childhood asthma and traffic flow using a geographic information system. *Environ Health Perspect.* 1999;107:761–767

Air Pollution and Children's Health

Joel Schwartz

Pediatrics 2004;113;1037

Updated Information & Services

including high resolution figures, can be found at:
http://pediatrics.aappublications.org/content/113/Supplement_3/1037

References

This article cites 81 articles, 12 of which you can access for free at:
http://pediatrics.aappublications.org/content/113/Supplement_3/1037#BIBL

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
Environmental Health
http://www.aappublications.org/cgi/collection/environmental_health_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



PEDIATRICS[®]

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Air Pollution and Children's Health

Joel Schwartz

Pediatrics 2004;113;1037

The online version of this article, along with updated information and services, is located on the World Wide Web at:

http://pediatrics.aappublications.org/content/113/Supplement_3/1037

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2004 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN[®]

