



LARC ICD-10 Coding Reference

Basic Implant Codes

ICD-10 Co	des – Implant	CPT Procedure Codes	HCPCSII/JCode	
Z30.017	Encounter for initial prescription of other contraceptives (includes Nexplanon) *updated 10/2016	11981 - insertion	J7307	
Z30.46	Encounter for surveillance of other contraceptives (includes removal, checking, reinsertion of Nexplanon) *updated 10/2016	11982 – removal 11983 – removal + reinsertion	J7307	
Z32.02	Pregnancy test/exam – negative	81025 (Urine pregnancy test)	Don't forget to bill for the point- of-care office pregnancy test (when conducted)	

Basic IUD Codes

ICD-10 Codes - IUDs		CPT Procedure Code	HCPCSII/JCode	
Z30.014	Encounter for initial prescription of IUD (not coded with the actual insertion)			
Z30.430	Encounter for insertion of IUD	58300	Kyleena = Q9984* Liletta = J7297 Mirena = J7298 ParaGard = J7300 Skyla = J7301	
Z30.431	Follow-up for patient with IUD or Routine checking for IUD			
Z30.432	Encounter for removal of IUD	58301		
Z30.433	Encounter for removal + reinsertion of IUD	58301 AND 58300-51* OR 58300-59* (Check with payer for expected modifier.) Append modifier -51 or -59 to the lesser paying service.	Kyleena = Q9984* Liletta = J7297 Mirena = J7298 ParaGard = J7300 Skyla = J7301	
Z32.02	Pregnancy test/exam – negative	81025 (Urine pregnancy test)	Don't forget to bill for the point- of-care office pregnancy test (when conducted)	





Coding Complex Cases (IUDs) with ICD-10

Clinical scenario	ICD-10 Code	s	CPT Procedure Code	HCPCSII/JCode
Failed insertion/ discontinued procedure	Z30.430 AND	Encounter for insertion of IUD	58300-52* or -53*	Kyleena = Q9984* Liletta = J7297
	<co-occurring complication which caused failed/ discontinued</co-occurring 	*Document reason for failed/stopped procedure with	*NOTE: Use modifier -52 (Failed Procedure) to denote that you attempted insertion but the procedure was incomplete due to anatomical factors (eg. Stenosis) or -53 (Discontinued Procedure) to indicate that you had to stop	Mirena = J7298 ParaGard = J7300 Skyla = J7301
	procedure>	appropriate ICD-10 codes.	because of concerns for patient well-being (eg. vaso-vagal, severe pain).	
Perforation (during insertion procedure)	Z30.430	Encounter for insertion of IUD	58300-53 *Use modifier - 53 if procedure was discontinued due to perforation.	Kyleena = Q9984* Liletta = J7297
	T83.39XA	Perforation of uterus by IUD (non-traumatic)		Mirena = J7298
				ParaGard = J7300 Skyla = J7301
Difficult insertion with ultrasound	Z30.430 AND	Encounter for insertion of IUD	58300-22 *Document the reason for additional work.	Kyleena = Q9984* Liletta = J7297
guidance	<co-occurring complication justifying</co-occurring 		76998 (Ultrasonic guidance, intraoperative) *Document the justification for ultrasonic guidance (eg. patient in severe pain).	Mirena = J7298 ParaGard = J7300
	ultrasound>			Skyla = J7301





Clinical scenario	ICD-10 Code	S	CPT Procedure Code	HCPCSII/JCode
Difficult insertion with ultrasound used to confirm the location of the IUD	Z30.430 <co-occurring complication justifying ultrasound></co-occurring 	Encounter for insertion of IUD *Document complication with appropriate ICD-10 codes.	 58300-22*Document the reason for additional work. 76857 Ultrasound, pelvic [nonobstetric], real time with image documentation; limited or follow-up -or - 76830 Ultrasound, transvaginal *NOTE: It is not routine practice to use ultrasound to confirm placement. You must document justification for ultrasonography (eg. Uterine perforation, severe pain). 	Kyleena = Q9984* Liletta = J7297 Mirena = J7298 ParaGard = J7300 Skyla = J7301
Missing strings, with ultrasound to locate	T83.32XA AND either Z30.431 OR Z30.432	Displacement of IUD - Missing strings, initial encounter	76857 Ultrasound, pelvic, limited or follow-up - OR - 76830 Ultrasound, transvaginal	
	Z30.431 Z30.432	Follow-up for patient with IUD (<i>if patient wants to keep IUD</i>) Encounter for IUD removal (<i>if patient desires removal</i>)	 58301 or 58301-22* *NOTE: You may append modifier -22 <i>if</i> the removal was complicated. Supporting documentation may be requested by a payer. If the string is easily located in the canal, -22 modifier should <i>not</i> be added. It should be appended only if it is a very difficult extraction and is separately documented with the claim. 	





Clinical scenario	ICD-10 Code	S	CPT Procedure Code	HCPCSII/JCode
Failed removal	Z30.432 AND	Encounter for IUD removal	58301-52 or -53*	
	T83.32XA OR	Displacement of IUD, initial encounter		
	<co-occurring complication which caused failed/ discontinued procedure></co-occurring 	*Document reason for failed/stopped procedure with appropriate ICD-10 codes.	*NOTE: Use modifier -52 to denote that you attempted removal but the removal procedure was incomplete (unable to remove/locate IUD) or modifier -53 to indicate that you had to stop because of concerns for patient well-being. You must document reason for failed or incomplete procedure.	





Common LARC Modifiers

Modifier	Definition	Possible Clinical Scenarios	Documentation
-22	Increased	Complex or difficult insertion	Total time of the procedure as compared with typical duration
	procedural	Unsuccessful insertion, followed by successful insertion	Reason for the additional work required
	services	during the same surgical session	Include diagnoses with appropriate ICD-10 codes or simple descriptive
			diagnoses that explain the reasons for the added difficulty
-25	Significant,	Patient came in for general contraceptive counseling,	Document time spent counseling
	separately	ends up choosing IUD or implant, and it is inserted that	
	identifiable E/M	day	
	service		
-51*	Multiple	• Removal of IUD and insertion of new IUD on the same day	Document should support the reasons for removal and reinsertion on
	procedures	Removal of implant and insertion of new implant on the	the same day (eg. IUD expired, desired to continue with same method)
	performed on the	same day	
	same day, during		
	the same session		
-52	Failed procedure	Provider couldn't complete procedure for anatomic	• Document reasons for procedure failure (eg. N88.2 Stricture/ stenosis
	-	reasons (eg. stenosis)	of cervix)
-53	Discontinued	Provider couldn't complete procedure due to concerns for	How much work was actually performed
	procedure	patient well-being	Document reason procedure terminated (eg. R55 Syncope/ vasovagal)
		Severe pain	
		Vasovagal	
		Patient changed mind during procedure	
-59*	Distinct		
-59*	procedural	Removal of IUD and insertion of new IUD on the same day	 Document should support the reasons for removal and reinsertion on the same day (or ULD outined desired to continue with same method)
	service	 Removal of implant and insertion of new implant on the same day. 	the same day (eg. IUD expired, desired to continue with same method)
		same day	
-76	Repeat procedure	Successful insertion but the IUD is expelled, followed by	 Document reason for repeat procedure (eg. IUD was expelled)
		repeat insertion	
L			

* When choosing between modifiers -51 and -59, payer policy may be the determining factor. Some payers will not pay for multiple procedures using modifier -51. *Check with payer*.