


Topic 11

Improving medication safety



Rationale

- Medication use has become increasingly complex in recent times
- Medication error is a major cause of preventable patient harm
- As future health-care workers, you will have an important role in making medication use safe

Learning objectives

- To provide an overview of medication safety
- To encourage students to continue to learn and practise ways to improve the safety of medication use

Knowledge requirements

- Understand the scale of medication error
- Understand the steps involved in a patient using medication
- Identify factors that contribute to medication error
- Learn how to make medication use safer
- Understand the benefits of a multidisciplinary approach to medication safety

Performance requirements

Acknowledge that medication safety is a topic and an understanding of the area will affect how you perform the following tasks:

- Use generic names where appropriate
- Tailor your prescribing for each patient
- Learn and practise thorough medication history taking
- Know which medications are high-risk and take precautions
- Be very familiar with the medication you prescribe and/or dispense
- Use memory aids
- Remember the 5 Rs when prescribing and administering
- Communicate clearly
- Develop checking habits
- Encourage patients to be actively involved in the process
- Report and learn from medication errors

Definitions (1)

- **Side-effect:** a known effect, other than that primarily intended, relating to the pharmacological properties of a medication
 - e.g. opiate analgesia often causes nausea
- **Adverse reaction:** unexpected harm arising from a justified action where the correct process was followed for the context in which the event occurred
 - e.g. an unexpected allergic reaction in a patient taking a medication for the first time
- **Error:** failure to carry out a planned action as intended or application of an incorrect plan
- **Adverse event:** an incident in which a patient is harmed

Source: Conceptual Framework for the International Classification for patient safety

Definitions (2)

- Adverse drug event:
 - May be preventable (e.g. the result of an error) or
 - May not be preventable (e.g. the result of an adverse drug reaction or side-effect)
- Medication error may result in ...
 - An adverse event if a patient is harmed
 - A near miss if a patient is nearly harmed or
 - Neither harm nor potential for harm
- Medication errors are preventable

Steps in using medication

- Prescribing
- Administering
- Monitoring

Note: these steps may be carried out by health-care workers or the patient; e.g. self-prescribing over-the-counter medication and self-administering medication at home

Prescribing involves ...

- Choosing an appropriate medication for a given clinical situation, taking individual patient factors into account, such as allergies
- Selecting the administration route, dose, time and regimen
- Communicating details of the plan with:
 - Whoever will administer the medication (written-transcribing and/or verbal)
 - And the patient
- Documentation

How can *prescribing* go wrong?

- Inadequate knowledge about drug indications and contraindications
- Not considering individual patient factors, such as allergies, pregnancy, co-morbidities, other medications
- Wrong patient, wrong dose, wrong time, wrong drug, wrong route
- Inadequate communication (written, verbal)
- Documentation - illegible, incomplete, ambiguous
- Mathematical error when calculating dosage
- Incorrect data entry when using computerized prescribing e.g. duplication, omission, wrong number

Look-a-like and sound-a-like medications

2 examples :

- Avanza (mirtazapine, antidepressant); Avandia (rosiglitazone, diabetes medicine)
- Celebrex (celecoxib, anti-inflammatory); Cerebryx (fosphenytoin, anticonvulsant); Celexa (Citalpram, antidepressant)

Ambiguous nomenclature

- Tegretol 100mg
- S/C
- 1.0 mg
- .1 mg
- Tegreto 1100 mg
- S/L
- 10 mg
- 1 mg

Avoiding ambiguous nomenclature

- Avoid trailing zeros
e.g. write 1 not 1.0
- Use leading zeros
e.g. write 0.1 not .1
- Know accepted local terminology
- Write neatly, print if necessary

Administration involves ...

- Obtaining the medication in a ready-to-use form; may involve counting, calculating, mixing, labeling or preparing in some way
- Checking for allergies
- Giving the right medication to the right patient, in the right dose, via the right route, at the right time
- Documentation

How can drug administration go wrong?

- Wrong patient
- Wrong route
- Wrong time
- Wrong dose
- Wrong drug
- Omission, failure to administer
- Inadequate documentation

The 5 Rs

- Right Drug
- Right Route
- Right Time
- Right Dose
- Right Patient

Calculation errors

Can you answer the following question?

A patient needs 300 micrograms of a medication that comes in a 1 ml ampoule containing 1 mg of the drug. What volume do you draw up and inject?

Monitoring involves ...

- Observing the patient to determine if the medication is working, being used appropriately and not harming the patient
- Documentation

How can monitoring go wrong?

- Lack of monitoring for side-effects
- Drug not ceased if not working, or course completed
- Drug ceased before course completed
- Drug levels not measured, or not followed up
- Communication failures

Do you know which drugs need blood tests to monitor levels?

Which patients are most at risk of medication error?

- Patients on multiple medications
- Patients with another condition, e.g. renal impairment, pregnancy
- Patients who cannot communicate well
- Patients who have more than one doctor
- Patients who do not take an active role in their own medication use
- Children and babies (dose calculations required)

In what situations are staff most likely to contribute to a medication error?

- Inexperience
- Rushing
- Doing two things at once
- Interruptions
- Fatigue, boredom, being on “automatic pilot” leading to failure to check and double-check
- Lack of checking and double checking habits
- Poor teamwork and/or communication between colleagues
- Reluctance to use memory aids

How can workplace design contribute to medication errors?

- Absence of a safety culture in the workplace
 - e.g. poor reporting systems and failure to learn from past near misses and adverse events
- Absence of memory aids for staff
- Inadequate staff numbers

How can medication presentation contribute to medication errors?

- Look-alike, sound-alike medications
- Ambiguous labeling

Ways to make medication use safer

What you can do to make medication use safer:

- Use generic names
- Tailor prescribing for individual patients
- Learn and practise collecting complete medication histories
- Know the high-risk medications and take precautions
- Be very familiar with the medications you prescribe
- Use memory aids
- Remember the 5 Rs
- Communicate clearly
- Develop checking habits
- Encourage patients to be actively involved
- Report and learn from errors

Use generic names rather than trade names

Tailor your prescribing for each individual patient

Consider:

- Allergies
- Co-morbidities (especially liver and renal impairment)
- Other medication
- Pregnancy and breastfeeding
- Size of patient

Learn and practise collecting complete medication histories

- Include name, dose, route, frequency, duration of every drug
- Ask about recently ceased medications
- Ask about over-the-counter medications, dietary supplements and complementary medicines
- Make sure what patient actually takes matches your list:
 - Be particularly careful across transitions of care
 - Practise medication reconciliation at admission to and discharge from hospital
- Look up any medications you are unfamiliar with
- Consider drug interactions, medications that can be ceased and medications that may be causing side-effects
- Always include allergy history

Know which medications are high risk and take precautions

- Narrow therapeutic window
- Multiple interactions with other medications
- Potent medications
- Complex dosage and monitoring schedules
- Examples:
 - Oral anticoagulants
 - Insulin
 - Chemotherapeutic agents
 - Neuromuscular blocking agents
 - Aminoglycoside antibiotics
 - Intravenous potassium
 - Emergency medications (potent and used in high pressure situations)

Be very familiar with the medications you prescribe

- Do some homework on every medication you prescribe
- Suggested framework
 - Pharmacology
 - Indications
 - Contraindications
 - Side-effects
 - Special precautions
 - Dose and administration
 - Regimen

Use memory aids

- Textbooks
- Personal digital assistant
- Computer programmes, computerized prescribing
- Protocols
- Free up your brain for problem solving rather than remembering facts and figures that can be stored elsewhere
- Looking things up if unsure is a marker of safe practice, not incompetence!

Remember the 5 Rs when prescribing and administering

Can you remember what they are?

- Right Drug
- Right Dose
- Right Route
- Right Time
- Right Patient

Communicate clearly

- State the obvious
- Write clearly and legibly
- The 5 Rs
- Close the loop

Develop checking habits (1)

- When prescribing a medication
- When administering medication:
 - Check for allergies
 - Check the 5 Rs
- Remember computerized systems still require checking
- Always check and it will become a habit!

Develop checking habits (2)

- Some useful maxims ...
- Unlabelled medications belong in the bin
- Never administer a medication unless you are 100% sure you know what it is
- Practice makes permanent, perfect practice makes perfect

Start your checking habits now !

Encourage patients to be actively involved in the process

- When prescribing a new medication provide patients with the following information:
 - Name, purpose and action of the medication
 - Dose, route and administration schedule
 - Special instructions, directions and precautions
 - Common side-effects and interactions
 - How the medication will be monitored
- Encourage patients to keep a written record of their medications and allergies
- Encourage patients to present this information whenever they consult a doctor

Report and learn from medication errors

Safe practice skills for students to develop and practise ...

- Whenever learning and practising skills that involve medication use, consider the potential hazards to the patient and what you can do to enhance patient safety
- Knowledge of medication safety will impact the way you:
 - Prescribe, document and administer medication
 - Use memory aids and perform drug calculations
 - Perform medication and allergy histories
 - Communicate with colleagues
 - Involve and educate patients about their medication
 - Learn from medication errors and near misses

Summary

- Medications can greatly improve health when used wisely and correctly
- Yet, medication error is common and is causing preventable human suffering and financial cost
- Remember that using medications to help patients is not a risk-free activity
- Know your responsibilities and work hard to make medication use safe for your patients

For discussion

- Are you aware of any incidents in which a patient was harmed by medication?
- Describe what happened
- Was the situation a result of a side-effect, adverse drug reaction or medication error?

Calculation errors (1)

- Can you answer the following question?

A 12 kg, 2-year-old boy requires 15 mg/kg of a medication that comes as a syrup with a concentration of 120 mg/5mls. How many mls do you prescribe?

Calculation errors (2)

- Can you answer the following question?

A patient needs 300 micrograms of a medication that comes in a 1 ml ampoule containing 1 mg of the drug. What volume do you draw up and inject?

Example case (1)

- A 74-year-old man sees a community doctor for treatment of new onset stable angina
- The doctor has not met this patient before and takes a full past history and medication history
- He discovers the patient has been healthy and only takes medication for headaches
- The patient cannot recall the name of the headache medication
- The doctor assumes it is an analgesic that the patient takes whenever he develops a headache



Example case (2)

- But the medication is actually a beta-blocker that he takes every day for migraine; this medication was prescribed by a different doctor
- The doctor commences the patient on aspirin and another beta-blocker for the angina
- After commencing the new medication, the patient develops bradycardia and postural hypotension
- Unfortunately the patient has a fall three days later due to dizziness on standing; he fractures his hip in the fall

What factors contributed to this medication error?

- Two drugs of the same class prescribed unknowingly with potentiation of side-effects
- Patient not well informed about his medications
- Patient did not bring medication list with him when consulting the doctor
- Doctor did not do a thorough enough medication history
- Two doctors prescribing for one patient
- Patient may not have been warned of potential side-effects and of what to do if side-effects occur

How could this situation have been prevented?

- Patient education regarding:
 - Regular medication
 - Potential side-effects
 - The importance of being actively involved in their own care - e.g. having a medication list
- More thorough medication history

Case (1)

- A 38-year-old woman comes to the hospital with 20 minutes of itchy red rash and facial swelling; she has a history of serious allergic reactions
- A nurse draws up 10 mls of 1:10,000 adrenaline (epinephrine) into a 10 ml syringe and leaves it at the bedside ready to use (1 mg in total) just in case the doctor requests it
- Meanwhile the doctor inserts an intravenous cannula
- The doctor sees the 10 ml syringe of clear fluid that the nurse has drawn up and assumes it is normal saline

Case (2)

- There is no communication between the doctor and the nurse at this time
- The doctor gives all 10 mls of adrenaline (epinephrine) through the intravenous cannula thinking he is using saline to flush the line.
- The patient suddenly feels terrible, anxious, becomes tachycardic and then becomes unconscious with no pulse
- She is discovered to be in ventricular tachycardia, is resuscitated and fortunately makes a good recovery
- Recommended dose of adrenaline (epinephrine) in anaphylaxis is 0.3 - 0.5 mg IM, this patient received 1mg IV

Can you identify the contributing factors to this error?

- Assumptions
- Lack of communication
- Inadequate labeling of syringe
- Giving a substance without checking and double-checking what it is
- Lack of care with a potent medication

How could this error have been prevented?

- Never give a medication unless you are sure you know what it is; be suspicious of unlabelled syringes
- Never use an unlabelled syringe unless you have drawn the medication up yourself
- Label all syringes
- Communication - nurse and doctor to keep each other informed of what they are doing
e.g. nurse: “I’m drawing up some adrenaline”
- Develop checking habits before administering every medication ... go through the 5 Rs
e.g. doctor: “What is in this syringe?”

Example case (1)

- A patient is commenced on oral anticoagulants in hospital for treatment of a deep venous thrombosis following an ankle fracture
- The intended treatment course is 3-6 months though neither the patient nor community doctor are aware of the planned duration of treatment
- Patient continues medication for several years, being unnecessarily exposed to the increased risk of bleeding associated with this medication

Example case (2)

- The patient is prescribed a course of antibiotics for a dental infection
- 9 days later the patient becomes unwell with back pain and hypotension, a result of a spontaneous retroperitoneal haemorrhage, requiring hospitalization and a blood transfusion
- International normalized ratio (INR) reading is grossly elevated, anticoagulant effect has been potentiated by the antibiotics

Can you identify the contributing factors for this medication error?

- Lack of communication and hence continuity of care between the hospital and the community
- Patient not informed of the plan to cease medication
- The interaction between antibiotic and anticoagulant was not anticipated by the doctor who prescribed the antibiotic even though this is a known phenomenon
- Lack of monitoring; blood tests would have detected the exaggerated anticoagulation effect in time to correct the problem

How could this error have been prevented?

- Effective communication
 - e.g. discharge letter from hospital to community doctor
 - e.g. patient information
- Memory aids and alerting systems to help doctors notice potential adverse drug interactions
- Being aware of common pitfalls in medications you prescribe
- Monitoring medication effects when indicated

How could the patient help prevent this error?

- Asking more questions:
 - “How long will I need this new medication for?”
 - “Will this antibiotic interact with my other medication?”
- How can the doctor encourage the patient to ask more questions?