

Dialysis and Sleep Problems

What are common sleep problems for people on Dialysis?

People on dialysis may complain of any the following problems:

- Difficulty falling asleep
- Trouble staying asleep
- Interrupted sleep
- Struggle to wake up

Why am I at Risk?

Studies show that up to 80% of dialysis patients have some form of sleep problems. Without treatment, sleep problems can lead to poor physical and mental functioning, lower quality of life, and increase death risk.

Why Might I Have Trouble Sleeping?

These may vary from person to person.

- Uremia -not getting enough dialysis.
- Daytime naps sleeping during the day makes it harder to sleep at night.
- Restless Legs Syndrome- the uncomfortable, "jumpy" feeling in your legs.
- Depression may cause you to sleep too much.
- Stress or anxiety worries may keep you awake at night.
- Itching may be the result of a high phosphorus level.
- Sleep Apnea- if you have difficulty breathing while sleeping you may need to be evaluated for sleep apnea.

What do I do if I have any of these symptoms?

Talk to your doctor and dialysis social worker if you are having trouble sleeping almost every night for more than two weeks. We can do a screening to evaluate the quality of your sleep and review treatment options with you.

What Are the Treatment Options for Sleep Problems?

- Your doctor may evaluate and treat your other medical conditions that impact your sleep.
- Your doctor may refer you to a sleep specialist for further evaluation.
- Your doctor may prescribe you a medication. Talk to your doctor before using over-the-counter sleep medications.
- Your dialysis social worker can meet with you to talk about your symptoms, provide support, and give you tips for sleeping better (see below).

What can I do to help myself Sleep Better?

- Go to bed and get up around the same time every day.
- Sleep in a dark, quiet room. Reserve the room for sleep and sex only.
- Drink less caffeine and maintain a regular meal schedule.
- Exercise regularly during the day.
- Avoid daytime naps.
- Do quiet activities until you are sleepy.
- Do not use nicotine.
- Avoid drinking alcohol before bed.
- Try a relaxation technique to reduce stress or worries

For More Information:

- University of Michigan Sleep Disorders Clinic: 734-936-9068
 - o https://medicine.umich.edu/dept/sleep/patient-information
- Life Options: <u>www.lifeoptions.org</u>

Disclaimer: This document contains information and/or instructional materials developed by the University of Michigan Health System (UMHS) for the typical patient with your condition. It may include links to online content that was not created by UMHS and for which UMHS does not assume responsibility. It does not replace medical advice from your health care provider because your experience may differ from that of the typical patient. Talk to your health care provider if you have any questions about this document, your condition or your treatment plan.

Author: Andrea DeKam, LMSW Reviewers: Karen Crampton, LMSW and Tracey Tyus-Bailey, LMSW

Patient Education by <u>University of Michigan Health System</u> is licensed under a <u>Creative Commons</u> <u>Attribution-NonCommercial-ShareAlike 3.0 Unported License</u>. Last Revised 05/2016